arXiv:1112.4515v1 [hep-ph] 19 Dec 2011

Unsolved problems in particle physics

Sergey Troitsky
Institute for Nuclear Research of the Russian Academy of Sciences

Abstract

I consider selected (most important according to my own choice) unsolved problems in particle theory, both those related to extensions of the Standard Model (neutrino oscillations, which probably do not fit the usual three-generation scheme; indications in favour of new physics from astrophysical observations; electroweak symmetry breaking and hierarchy of parameters) and those which appear in the Standard Model (description of strong interactions at low and intermediate energies).

Contents

1 Introduction: status and parameters of the Standard Model 2

2 The observed deviation from the Standard Model: neutrino oscillations. 5
2.1 Theoretical description. 6
2.2 Experimental results: standard three-flavour oscillations. 7
2.3 Experimental results: non-standard oscillations. 10
2.4 The neutrino mass. 13

3 Astrophysical and cosmological indications in favour of new physics. 13
3.1 Baryon asymmetry. 14
3.2 Dark matter. 15
3.3 Accelerated expansion of the Universe 19

4 Aesthetic difficulties: the origin of parameters. 23
4.1 Electroweak interaction and the Higgs boson. 23
4.2 The gauge hierarchy. 26
4.3 The fermion mass hierarchy. 35

5 Theoretical challenges in the description of hadrons. 37
5.1 Problems of the perturbative QCD. 37
5.2 The lattice results. 39
5.3 Dual theories: supersymmetric duality and holography. 40

6 Conclusions. 43

1 Introduction: status and parameters of the Standard Model

One may compare the current state of quantum field theory and its applications to particle physics with the situation 20-30 years ago and discover, amusingly, that all principal statements of this field of physics are practically unchanged, which is in contrast with rapid progress in condensed-matter physics. Indeed, most of the experiments, held during the last two decades, supported the correctness of predictions which had been made earlier, derived from the models developed earlier. This success of the particle theory resulted in considerable stagnation in its development. However, one may expect that in the next few years, the particle physics will again become an intensively developing area. Firstly, there is a certain amount of collected experimental results (first of all related to cosmology and astrophysics, but also obtained in laboratories) which suggest that the Standard Model (SM) is incomplete. Secondly, the theory was developing under the guidance of the principle of naturalness, that is the requirement to explain quantitatively any hierarchy in model parameters (in the case of SM, it is possible only within a larger fundamental theory yet to be constructed). Finally, one of the most important arguments for the coming excitement in particle physics is the expectation of new results from the Large Hadron Collider. As it will become clear soon, this accelerator will be able to study the *full* range of energies where the physics responsible for the electroweak symmetry breaking should appear, so we are expecting interesting discoveries in the next few years in any case: either it will be the Higgs boson, or some other new particles, or (in the most interesting case) no new particle will be found which would suggest a serious reconsideration of the Standard Model.

The Large Hadron Collider (LHC, see e.g. [1]) is an accelerator which allows to collide protons with the center-of-mass energy up to 14 TeV (currently working at 7 TeV) and heavy nuclei. In a 30-km length tunnel, at the border of Switzerland and France, there are four main experimental installations (general-purpose detectors ATLAS and CMS; LHCb which is oriented to the study of B mesons and ALICE, specialized in heavy-ion physics) as well as a few smaller experiments. The first results of the work of the collider have brought a lot of new information on particle interactions which we will mention when necessary.

The purpose of the present review is to discuss briefly the current state of particle physics and possible prospects for its development. For such a wide subject, the selection of topics is necessarily subjective and estimates of importance of particular problems and of potential of particular approaches reflect the author's personal opinion, while the bibliography cannot be made exhaustive.

The contemporary situation in the particle physics may be described as follows. Most of the modern experimental data are well described by the Standard Model of particle physics which was created in 1970s. At the same time, there are a considerable amount of indications that SM is not complete and is not more than a good approximation to the correct description of particles and interactions. We are not speaking now about minor deviations of certain measured observables from theoretically calculated ones – these deviations may be related to insufficient precision of either the measurement or the calculations, to unaccounted systematic errors or insufficient sets of experimental data (statistical fluctuations); it happens that these deviations disappear after a few years of more detailed study. Contrary, we will emphasise more serious qualitative problems of SM, the latter being considered as an instrument of quantitative description of elementary particles. These problems include the following:



Figure 1: Particle interactions.

(1) experimental indications to the incompleteness of SM, namely the well-established experimental observations of neutrino oscillations (which are impossible, see Sec. 2.4, in SM) and incapability of SM to describe the results of astrophysical observations, in particular of those related to the structure and evolution of the Universe;

(2) not fully natural and not calculable in the theory values of the SM parameters, in particular, the fermion mass hierarchy, the hierarchy of symmetry-breaking scales and the absence of a light (with mass $\lesssim 100$ GeV) Higgs boson;

(3) purely theoretical difficulties in description of hadrons by means of the available methods of quantum field theory.

We will discuss these unsolved problems of SM and related prospects for the development of the particle theory.

For future reference, it is useful to recall briefly the structure of SM (see e.g. [2, 3] and the appendix to [4]). The model includes a certain set of particles and their interactions.

Out of four known interactions (see Fig. 1), three are described by SM – the electromagnetic, weak and strong ones. The first two of them have a common *electroweak* gauge interaction behind them. The symmetry of this interaction, $SU(2)_{\rm L} \times U(1)_{\rm Y}$, manifests itself at energies higher than ~ 200 GeV. At lower energies, this symmetry is broken down to $U(1)_{\rm EM} \neq U(1)_{\rm Y}$ (the electroweak symmetry breaking); in SM, this breaking is related to the vacuum expectation value of a scalar field, the Higgs boson. Parameters of the electroweak breaking are known up to a high precision; experimental data are in a perfect agreement with the theory. The Higgs boson has not been observed yet; its mass, being a free parameter of the theory, is bound by direct experimental searches (see Table 1 and more details in Sec. 4.1).

The *strong* interaction in SM is described by the quantum chromodynamics (QCD), a theory with the gauge group $SU(3)_{\rm C}$. The effective coupling constant of this theory grows when the energy is decreased. As a result, particles which feel this interaction cannot exist as free states and appear only in the form of bound states called hadrons. Most of modern methods of quantum field theory work for small values of coupling constants, that is, for QCD, at high energies.

The fourth known interaction, the *gravitational* one, is not described by SM, but its effect on the microscopic physics is negligible.

The particle content of SM is summarized in Fig. 2. Quarks and leptons, the so-called SM matter fields, are described by fermionic fields. Quarks take part in strong interactions and compose observable bound states, hadrons. Both quarks and leptons participate in the electroweak interaction. The matter fields constitute three generations; particles from different generations interact identically but have different masses. The full electroweak symmetry forbids fermion masses, so nonzero masses of



Figure 2: Particles described by the Standard Model.

quarks and leptons are directly related to the electroweak breaking – in SM, they appear due to the Yukava interaction with the Higgs field and are proportional to the vacuum expectation value of the latter. For the case of neutrino, these Yukawa interactions are forbidden as well, so neutrinos are strictly massless in SM. The gauge bosons, which are carriers of interactions, are massless for unbroken gauge groups $U(1)_{\rm EM}$ (electromagnetism – photons) and $SU(3)_{\rm C}$ (QCD – gluons); masses of $W^\pm$ and Z bosons are determined by the mechanism of the electroweak symmetry breaking. All SM particles, except for the Higgs boson, are found experimentally.

From the quantum-field-theory point of view, quarks and leptons may be described as states with definite mass. At the same time, gauge bosons interact with superpositions of these states; in another formulation, when the base is chosen to consist of the states interacting with the gauge bosons, the SM symmetries allow not only for the mass terms, $m_{ii}\bar\psi_i\psi_i$, for each ith fermion ψ_i, but also for a nondiagonal *mass matrix* $m_{ij}\bar\psi_i\psi_j$. Up to unphysical parameters, in SM, this matrix is trivial in the leptonic sector, while in the quark sector it is related to the Cabibbo-Kobayasi-Maskava (CKM) matrix. The latter may be expressed through three independent real parameters (quark mixing angles) and one complex phase (for more details, see [3, 5]).

The Standard Model has therefore 19 independent parameters, values of 18 of which are determined experimentally. They include three gauge coupling constants, α_s, α_2 and α_1 for gauge groups $SU(3)_C$, $SU(2)_W$ and $U(1)_Y$, respectively (the latter two are often expressed through the electromagnetic coupling constant α and the mixing angle θ_W), the QCD Θ-parameter, nine charged-fermion masses $m_{u,d,s,c,b,t,e,\mu,\tau}$, three quark mixing angles $\theta_{12,13,23}$, one complex phase δ of the CKM matrix and two parameters of the Higgs sector, which are conveniently expressed through the known Higgs-boson vacuum expectation value v and its unknown mass M_H. Experimental values of these parameters, recalculated from the 2010 data [6] (bounds on the mass of the Higgs boson based on LEP, Tevatron and LHC data are given as of December, 2011), may be found in Table 1.

It is worth reminding that the observable world is mostly mad eof atoms, so, out of

$\alpha_s(M_Z)$	=	0.114 ± 0.0007
$1/\alpha(M_z)$	=	127.916 ± 0.015
$\sin^2\theta_W(M_Z)$	=	0.23108 ± 0.00005
Θ	$\lesssim$	10^{-10}
$m_u(2\text{ GeV})$	=	$2.5^{+0.8}_{-1.0}$ MeV
$m_d(2\text{ GeV})$	=	$5.0^{+1.0}_{-1.5}$ MeV
$m_s(2\text{ GeV})$	=	105^{+25}_{-35} MeV
$m_c(m_c)$	=	$1.266^{+0.031}_{-0.036}$ GeV
$m_b(m_b)$	=	4.198 ± 0.023 GeV
$m_t(m_t)$	=	173.1 ± 1.35 GeV
m_e	=	510.998910 ± 0.000013 keV
m_μ	=	105.658367 ± 0.000004 MeV
m_τ	=	1.77682 ± 0.00016 GeV
θ_{12}	=	$13.02^\circ \pm 0.05^\circ$
θ_{23}	=	$2.35^\circ \pm 0.06^\circ$
θ_{13}	=	$0.199^\circ \pm 0.011^\circ$
δ	=	1.20 ± 0.08
$v(m_\mu)$	=	246.221 ± 0.002 GeV
m_H		115 GeV ... 127 GeV

Table 1: Parameters of the Standard Model. For parameters with significant energy dependence, the energy scales, to which the numerical values correspond, are given in parentheses.

the full manifold of elementary particles, only few are met "in the everyday life". These are u and d quarks in the form of protons (udd) and neutrons (uud), electrons and, out of interaction carriers, the photon. The reasons for that are different for different particles. In particular, neutrino does not interact with the electromagnetic field and is therefore very hard to detect; heavy particles are unstable and decay to lighter ones; strongly interacting quarks and gluons are confined in hadrons. The full manyfold of SM particles reveal themselves either in complicated dedicated experiments, or indirectly by their effect seen in astrophysical observations.

Thus, before to proceed with the description of unsolved problems, let us recall that all experimental results concerning physics of charged leptons, photons, W and Z bosons at all available energies and quarks and gluons at high energies are in excellent agreement with SM for a given set of its parameters.

2 The observed deviation from the Standard Model: neutrino oscillations.

Let us discuss the unique, well-established in laboratory experiments, evidence in favour of incomleteness of SM, the phenomenon of neutrino oscillations, that is mutual conversion of neutrinos of different generations to each other. A more detailed modern description of the problem may be found in the book [7], in the Appendix to the textbook [4], in reviews [8, 9, 10] etc.

2.1 Theoretical description.

In analogy with the case of charged leptons, let us consider three generations of neutrino: electron neutrino (ν_e), muon neutrino (ν_μ) and tau neutrino (ν_τ). The corresponding fermion fields interact with the gauge bosons W and Z through weak charged and neutral currents. These interactions are responsible for both creation and experimental detection of neutrinos.

Similarly to the quark case, one may suppose that neutrinos have a nonzero mass matrix (though it cannot be incorporated in SM, the low-energy effective theory, electrodynamics, does not forbid it) which may be nondiagonal. It is convenient to describe this system in terms of linear combinations $\nu_{1,2,3}$ of the original fields $\nu_{e,\mu,\tau}$ with the diagonal mass matrix,

$$\nu_i = \sum_{\alpha=e,\mu,\tau} U_{i\alpha}\nu_\alpha,$$

where $U_{i\alpha}$, $i = 1, 2, 3$; $\alpha = e, \mu, \tau$, are the elements of the leptonic mixing matrix.

To demonstrate the phenomenon of neutrino oscillations, let us restrict ourselves to the case of two flavours, ν_e and ν_μ. Let their linear combinations,

$$\nu_1 = \cos\theta_{12}\,\nu_e + \sin\theta_{12}\,\nu_\mu, \tag{1}$$

$$\nu_2 = -\sin\theta_{12}\,\nu_e + \cos\theta_{12}\,\nu_\mu,$$

be the eigenvectors of the mass matrix with eigenvalues m_1^2, m_2^2, respectively. The inverse transformation expresses (ν_e, ν_μ) through (ν_1, ν_2):

$$\nu_e = \cos\theta_{12}\,\nu_1 - \sin\theta_{12}\,\nu_2,$$

$$\nu_\mu = \sin\theta_{12}\,\nu_1 + \cos\theta_{12}\,\nu_2.$$

Suppose that at the moment $t = 0$, in a certain weak-interaction event, an electron neutrino ν_e was created, that is the superposition of ν_1 and ν_2 with known coefficients:

$$\nu_1(0) = \cos\theta_{12}\,\nu_e(0),$$

$$\nu_2(0) = -\sin\theta_{12}\,\nu_e(0).$$

The evolution of mass eigenstates for a plane monochromatic wave moving in the direction z mas be described as

$$\nu_i(z,t) = \mathrm{e}^{-i\omega t + i\sqrt{\omega^2 - m_i^2}z}\nu_i(0), \quad i = 1, 2,$$

where ω is the energy and $\sqrt{\omega^2 - m_i^2}$ is the momentum. While propagating, the wave packets corresponding to ν_1 and ν_2 disperse in different ways, so that the relation $(\cos\theta_{12}, -\sin\theta_{12})$ between their coefficient no longer holds, which means that an admixture of the orthogonal state, ν_μ, appears. In the (commonly considered) ultrarelativistic limit, $\omega \gg m_i$ and $\sqrt{\omega^2 - m_i^2} \simeq \omega - \frac{m_i^2}{2\omega}$. The probability to detect ν_μ at a point (t, z) for each emitted ν_e is then

$$P(\nu_\mu; z, t) = |\nu_\mu(z,t)|^2 = \sin^2 2\theta_{12}\,\sin^2\left(\frac{m_2^2 - m_1^2}{4\omega}z\right). \tag{2}$$

One may see that this probability is an oscillating function of the distance z, hence the term "neutrino oscillations". As expected, no oscillations happen either in the case of equal (even nonzero) masses (similar dispersions of ν_1 and ν_2) or for a diagonal mass matrix ($\theta_{12} = 0$, $\nu_1 = \nu_e$ etc.). A similar description of oscillations of three neutrino flavours determines, in analogy with Eq. (1), three mixing angles θ_{12}, θ_{13}, θ_{23}.

When individual neutrinos propagate to large distances, the oscillation formalism described above stops to work, because the particles of different mass require different time to propagate from the source, hence loss of coherence; nevertheless the transformations of neutrinos are possible and their probability is calculable.

2.2 Experimental results: standard three-flavour oscillations.

Let us turn now to the history (see e.g. [7]) and the modern state (see e.g. [11]) of the question of neutrino oscillations. In 1957, Pontecorvo [12, 13] suggested the possibility of oscillations in the "neutrino–antineutrino" system, similar to K meson oscillations already known at that time. This first mentioning of the possibility of neutrino oscillations was aimed at the explanation of preliminary Davis' results about observation of the reaction $\bar{\nu} + {}^{37}\mathrm{Cl} \to {}^{37}\mathrm{Ar} + e^-$ with reactor neutrinos. On one hand, this experimental result has not been confirmed; on the other one, it has become clear that the Pontecorvo model was not able to describe it even if it were true. The first mention of mutual transformations of ν_e and ν_μ is due to Maki, Nakagawa and Sakata [14], while the first succesful description of oscillations in the system of two-flavour neutrinos was given by Pontecorvo [15] and by Gribov and Pontecorvo [16]. The theory of neutrino oscillations in its present form has been developed in 1975-76 by Bilenky and Pontecorvo [17, 18], Eliser and Swift [19], Fritch and Minkowski [20], Mikheyev, Smirnov [21, 22] and Wolfenstein [23].

The first experimental evidence in favour of neutrino oscillations have been obtained more than a half century ago, though for considerable period of time their interpretation remained an open question. We are speaking about the so-called "solar neutrino problem": the observed flux of neutrinos form the Sun was considerably lower than it was predicted by a model of solar nuclear reactions. This solar neutrino deficit was first found in the Homestake experiment (USA; first as early as 1968 [24]) and subsequently confirmed by Kamiokande (Japan) [25], SAGE (Russia, Baksan neutrino observatory of INR, RAS) [26], GALLEX/GNO (Italy, the Gran-Sasso laboratory) [27] and Super-K (Japan) [28] experiments, which made use of various experimental techniques and were sensitive to neutrinos form different nuclear reactions. Since only electron neutrinos are produced in the Sun, and only these were detected in the experiments, the deficit might be explained by transformation of a part of electron neutrinos to muon ones.

The natural source of *muon* neutrinos is provided by cosmic rays, that is charged particles (protons and nuclei) of extraterrestrial origin which interact with atoms in the Earth's atmosphere and produce secondary particles. A significant part of the latters are charged π mesons. Neutrinos from decays of these π mesons, as well as from decays of secondary muons, are called atmospheric neutrinos. The first indications to oscillations of the atmospheric neutrinos have been obtained in the end of 1980s in Kamiokande [29] and IMB [30] experiments, with subsequent confirmation in Soudan-2 [31], MACRO [32] and Super-K [33]. Their result is the anisotropy in the flux of muon neutrinos: from above, that is from the atmosphere, the flux is higher than from below (through the Earth). Without oscillations, the flux were isotropic since it is determined by an isotropic flux of primary cosmic rays while the interaction of neutrino with the terrestrial matter is negligible. This anisotropy is not seen for electron neutrinos, hence



Figure 3: Limits (95% confidence level) on the $\nu_e - \nu_\mu$ oscillation parameters as result from the analysis taking into account three neutrino flavours [36]. The dotted line corresponds to the combination of solar experiments, the full line represents the KamLAND constraints, the gray ellipsis gives the constraints from the combination of all data. The star, the triangle and the square correspond to the most probable oscillation parameters obtained in these three analyses, respectively.

it is natural to suppose that ν_μ oscillate mainly to ν_τ (the latters were not detected in these experiments).

In the first decade of our century, a significant experimental progress in the questions we discuss has been achieved, so that now we have a reliable experimental proof of neutrino transformations with measured parameters.

$\nu_e - \nu_\mu$ **oscillations.** In addition to more or less model-dependent results about the solar neutrino deficit (ν_e *disappearence*), the SNO experiment has detected, in 2001 [34], *appearence* of neutrino of other flavours from the Sun in a full agreement with the flux expected in the oscillational picture. It has therefore closed the "solar neutrino problem" and supported, at the same time, the standard solar model. The KamLAND experiment [35] registered disappearence of electron antineutrino born in reactors of atomic power plants (in contrast with the case of the Sun, the initial flux of the particles may be directly determined in this case). Parameters of oscillations, measured in these very different experiments, are in excellent agreement, see Fig. 3. The SNO results, together with even more precise results of the BOREXINO experiment (Italy) [37], confirm the expected energy dependence of the number of disappeared solar neutrinos in agreement with predictions of Mikheyev, Smirnov [21, 22] and Wolfenstain [23], who developed a theory of neutrino oscillations in plasma: due to the fact that electrons are present in plasma, unlike muons and tau leptons, the interaction with medium goes differently for different types of neutrino. As a result, the oscillation formalism is modified and the resonance enhancement of oscillations becomes possible.

$\nu_\mu - \nu_\tau$ **oscillations.** In addition to the Super-K experiment, which have measured [38, 39] deviations from isotropy in atmospheric ν_μ and $\bar{\nu}_\mu$ to a great accuracy, the disappearence of ν_μ has been measured directly in neutrino beams created by particle accelerators (experiments K2K [40] and MINOS [41]), see Fig. 4. Finally, in 2010, the OPERA detector which is located in the Gran Sasso laboratory (Italy) has detected



Figure 4: Limits (90% confidence level) on the $\nu_\mu - \nu_\tau$ oscillation parameters. The dotted line represents the results of the SuperK analysis with account of three neutrino flavours [39]; the full line represents constraints by MINOS [42]. The star and the triangle denote the most probable oscillation parameters for these two analyses, correspondingly.

Δm^2_{12}	=	$\left(7.58^{+0.22}_{-0.26}\right) \times 10^{-5}$ eV^2
Δm^2_{23}	=	$\left(2.31^{+0.12}_{-0.09}\right) \times 10^{-3}$ eV^2
$\sin^2\theta_{12}$	=	$0.312^{+0.017}_{-0.016}$
$\sin^2\theta_{13}$	=	0.025 ± 0.007
$\sin^2\theta_{23}$	=	$0.42^{+0.08}_{-0.03}$

Table 2: Parameters of oscillations of three flavours of neutrino obtained with account of all relevant experimental data as of summer 2011 [47].

[43] the first (and currently unique) case of *appearence* of ν_τ in the ν_μ beam from the SPS accelerator (CERN, Switzerland).

The mixing angle θ_{13}. For a long time, the solar $(\nu_e - \nu_\mu)$ and atmospheric $(\nu_\mu - \nu_\tau)$ oscillation data have been described independently (see discussions in [4], Appendix C) while relatively low precision of experiments allowed for zero value of the mixing angle θ_{13}. The situation has been recently changed and, analyzed commonly, the data of various experiments point to nonzero θ_{13} [44]. In summer 2011, two accelerator experiments, T2K (Japan) [45] and MINOS [46], which both search for appearence of ν_e in ν_μ beams, published their results which are incompatible with $\theta_{13} = 0$. A quantitative analysis of all data on the solar and atmospheric neutrinos, jointly with the accelerator and reactor experiments which study the same part of the parameter space, points [47] towards a nonzero value of θ_{13} at the confidence level better than 99%. In Table 2, the results of this analysis are quoted.

2.3 Experimental results: non-standard oscillations.

The combination of all experiments described above is in a good quantitative agreement with the picture of oscillations of three types of neutrino with certain parameters. However, there exist results which do not fit this picture and may suggest that the fourth (or, maybe, even the fifth) neutrino exists. As we have seen above, one of the principal oscillation parameters is the mass-square difference, $\Delta m^2_{ij} = m^2_j - m^2_i$. The results on atmospheric and solar neutrinos, jointly with the accelerator and reactor experiments, are explained by two unequal Δm^2_{ij}, see Table 2,

$$\Delta m^2_{12} \ll \Delta m^2_{23} \sim 2 \times 10^{-3} \text{ eV}^2.$$

In the case of three neutrinos, these two values compose the set of linearly independent Δm^2_{ij} and

$$|\Delta m^2_{13}| = |\Delta m^2_{12} - \Delta m^2_{23}| \sim \Delta m^2_{23}.$$

Therefore, the observation of any neutrino oscillations with $\Delta m^2_{ij} \gg \Delta m^2_{23}$ implies either the existence of a new neutrino flavour ($i, j > 3$) or some other deviation from the standard picture. On the other hand, there is a very restrictive bound on the number of relatively light ($m_i < M_Z/2$) particles with the quantum numbers of neutrino. This bound comes from precise measurements of the Z-boson width and implies that there are only three such neutrinos. This means that the fourth neutrino, if exists, does not interact with the Z boson; in other words, it is "sterile". We will turn now to a certain experimental evidence in favour of $\Delta m^2_{ij} \sim 1$ eV^2. We note that the oscillations related to this Δm^2_{ij} should reveal themselves at relatively short distances and may be detected in so-called short-baseline experiments.

$\bar{\nu}_\mu - \bar{\nu}_e$ **oscillations.** The LSND experiment [48] studied muon decay at rest, $\mu^+ \to e^+ \nu_e \bar{\nu}_\mu$, and measured the $\bar{\nu}_e$ flux at the distance about 30 m from the place where muons were held. The excess of this flux over the background rate has been detected and interpreted as appearence of $\bar{\nu}_e$ as a result of $\bar{\nu}_\mu$ oscillations, for a range of possible parameters. A similar experiment, KARMEN [49] excluded a significant part of this parameter space, however, in 2010, the MiniBooNE experiment [50] has also detected an anomaly which is compatible with the LSND results and, within statistical uncertainties, does not contradict KARMEN for a certain range of parameters (Fig 5).

Another group of short-baseline experiments which study possible $\bar{\nu}_e - \bar{\nu}_\mu$ oscillations search for disappearence of $\bar{\nu}_e$ in the antineutrino flux from nuclear reactors. These experiments continued for decades; recently, their results have been reanalized jointly [52] with a more precise theoretical calculation of the expected fluxes. It has been shown that there is a statistically significant deficit of $\bar{\nu}_e$ in the detectors which is compatible with $\Delta m^2 \sim 1\text{eV}^2$, – the so-called reactor neutrino anomaly. The corresponding limits on the parameters are also shown in Fig. 5 for convenience. However, one should keep in mind that while LSND, KARMEN and MiniBooNE detected $\bar{\nu}_e$ in the $\bar{\nu}_\mu$ flux, therefore constraining $\bar{\nu}_e - \bar{\nu}_\mu$ oscillations, the reactor experiments only fix the disappearance of $\bar{\nu}_e$. While the lack of this disappearence excludes $\bar{\nu}_e - \bar{\nu}_\mu$ oscillations, the presence of it may be explained as a transformation of $\bar{\nu}_e$ into antineutrino of any other type.

One can see that there are several independent indications in favour of $\Delta m^2 \sim 1\text{eV}^2$, which, as we discussed above, require either introduction of more-than-three neutrino flavours or (see below) some other new physics.



Figure 5: Limits (90% confidence level) on the parameters of $\bar{\nu}_\mu - \bar{\nu}_e$ oscillations. The shaded region is compatible with the LSND signal [48]; the region inside the dotted curve – with the MiniBooNE signal [51]. Thin full lines bound the region of parameters compatible with a joint re-analysis of reactor data [52], see text. The KARMEN2 experiment excludes [49] the region above and to the right from the thick full line.

Other anomalies. Recent intense exploration of the field of neutrino oscillations revealed also a range of other anomalies which are currently being discussed and re-checked intensively.

Possible difference between neutrino and antineutrino oscillations. The MiniBooNE experiment studied separately neutrino and antineutrino beams. The $\bar{\nu}_e$ appearence has been detected [50, 51] while that of ν_e has not [53] (see Fig. 6). In assumption of equal oscillation parameters for ν and $\bar{\nu}$, the MiniBooNE result contradicts to LSND, but without this assumption, contrary, the LSND claim is supported. It is worth noting that the MINOS experiment also performed separate measurements with neutrino and antineutrino beams (studying a range of much smaller Δm^2); first their results for the two cases were incompatible at the 98% confidence level, however, subsequent analysis of a larger amount of data did not confirm this difference [54]. The latter result agrees with Super-K: though this experiment cannot distinguish neutrino from antineutrino in each particular case, it may limit [55] antineutrino oscillation parameters statistically, on the basis of a known contribution of $\bar{\nu}_\mu$ to the atmospheric neutrino flux.

Calibration of gallium detectors. The GALLEX [56] and SAGE [57] experiments, constructed to detect solar neutrinos with the help of the gallium detectors, calibrated their instruments with the help of artificial sources of radioactivity. They detected a deficit of electron neutrino compatible with oscillations with $\Delta m^2 \gtrsim 0.1$ eV^2 (see also [58]). This mass-square difference, which by itself does not agree with the standard three-neutrino oscillation picture, agrees with the antineutrino results of LSND, MiniBooNE and reactor experiments, however the corresponding mixing angle differs from the predictions of the latters [59].

Other puzzles. When speaking about unexplained results of neutrino experiments, one may mention also the unexpected excess of events with energies $\lesssim 400$ MeV detected by MiniBooNE for neutrinos [60] and antineutrinos [51]; possible seasonal variations of the neutrino flux in the Troitsk-νmass [61] and MiniBooNE [62] experiments; the result of the OPERA experiment [63] which measured the velocity of muon neutrinos



Figure 6: Limits (90% confidence level) on the $\nu_\mu - \nu_e$ and $\bar{\nu}_\mu - \bar{\nu}_e$ oscillation parameters. The shaded area corresponds to the part of the parameter space which is excluded for neutrino oscillations by MiniBooNE [53] and KARMEN [49], while thick contours limit the region which corresponds to the signal in antineutrino oscillations (full lines, MiniBooNE [51]; dotted line, LSND [48]).

which happened to be large than the speed of light. All these very interesting anomalies currently await their confirmation in independent experiments.

Possible theoretical explanations. The experimental results listed above are rather hard to explain. On one hand, a series of experiments suggest neutrino transformations compatible with $\Delta m^2 \gtrsim 0.1$ eV^2 which cannot be described in the frameworks of a standard three-generation scheme. On the other hand, the addition of the fourth neutrino cannot help to explain the difference between the neutrino and antineutrino oscillations [64, 65]. Alternatively, one can consider (a) two generations of sterile neutrinos (see e.g. [66] and references therein); (b) breaking [67] of the CPT invariance[1], or (c) nonstandard interaction of neutrino with matter which may distinguish particles and antiparticles [71, 72]. A critical analysis of some of these suggestions may be found e.g. in [66, 73, 74]. These scenarios experience considerale difficulties with simultaneous explanation of the full set of the experimental data, though they cannot be totally excluded; it might happen that a certain combination of these possibilities is realized in Nature.

A confirmation of the result about superluminal neutrino motion would require a serious reconsideration of basic ideas of particle physics. A successful theory which explains quantitatively the OPERA result should also agree with very restrictive bounds on the Lorentz-invariance violation in the sector of charged particles, with the absence

[1] Invariance with respect to simultaneous charge conjugation (C) and reflection of both space (P) and time (T) coordinates is (see e.g. [68]) a fundamental symmetry which inevitably exists in any (3+1)-dimensional Lorentz-invariant local quantum field theory. However, there exist phenomenologically acceptable models with the CPT violation (either with higher number of space dimensions, or with the Lorentz invariance violation, or with a nonlocal interaction). In the context of the neutrino oscillations, they are discussed e.g. in [67, 69, 70].

of dispersion of the neutrino signal from the supernova 1987A and with absence of intense neutrino decays which are characteristic for many models with deviations from the relativistic invariance.

2.4 The neutrino mass.

Conversions of neutrino of one type to another are experimentally proven and the set of numerous independent and very different experiments are in a good agreement with the oscillation picture. The oscillatory behaviour of the neutrino conversions is proven by a comparison of the results obtained for different energies (cf. the argument of the sine squared in Eq. (2)). The last step is to measure the neutrino flux at different distances along a single path (the distance dependence in Eq. (2)) which is planned for the nearest future. Up to this last detail, the neutrino oscillations are experimentally confirmed. Since the oscillations are possible only for different masses of neutrinos of different types, they prove also that (at least some of) the neutrino masses are nonzero. At the same time, direct experimental searches for neutrino masses have not been succesful yet; the most restrictive bounds, put by the Troitsk-νmass (INR RAS) and Mainz experiments, where the tritium beta decay was studied, are $m_{\nu_e} \lesssim 2$ eV [75, 76]. For other neutrino types, the experimental bounds on the mass are much weaker. An indirect bound on the sum of the neutrino masses may be obtained from the studies of anisotropies of the cosmic microwave background and of hierarchy of structures in the Universe [77]; it reads $\sum_i m_{\nu_i} \lesssim 0.35$ eV.

At the same time, in SM the lepton numbers are conserved separately for each generations, that is changes of the neutrino flavour are forbidden. By making use of the SM fields, it is impossible to construct a gauge invariant renormalizable interaction resulting in the neutrino mass, even after the electroweak symmetry breaking. Therefore, neutrino oscillations represent an experimental proof of the incompleteness of SM.

How can one modify SM to have massive neutrinos? First note that at energies below the electroweak breaking scale, the neutrino field is gauge invariant, it is uncharged and colorless. For such fermion fields one may write two kinds of mass terms, namely the Dirac term $m_{\mathrm{D}}\bar{\nu}_{\mathrm{R}}\nu_{\mathrm{L}}$ (all charged SM fermions have similar masses) and the Majorana one, $m_{\mathrm{M}}\nu_{\mathrm{L}}C\nu_{\mathrm{L}}$, where C is the charge conjugation matrix and ν_{L}, ν_{R} denote the left-handed and right-handed neutrino spinors, respectively.

In SM, only left-handed neutrinos are present, therefore to have Dirac masses, one must introduce new fields $\nu_{\mathrm{R},i}$. At first sight, the Majorana mass does not require new fields; however, like the Dirac one, it cannot be obtained from a renormalizable interaction. Going beyond the renormalizability means that SM is a low-energy limit of a more complete theory (like the non-renormalizable Fermi theory is a low-energy limit of SM), so it is again inevitable to introduce new fields. In any case, neutrinos are several orders of magnitude lighter than the charged fermions and a succesful theory of neutrino masses should be able to explain this fact (see also Sec. 4.3).

3 Astrophysical and cosmological indications in favour of new physics.

While laboratory experiments in particle physics give only limited indications to the incompleteness of SM (neutrino oscillations being the main one), most scientists are confident that a more complete theory should be invented. The main reason for this confidence comes from astrophysics and cosmology. In recent decades, intense devel-

opment of the observational astronomy in various energy bands has forced cosmology (that is the branch of science studying the Universe as a whole) to become an accurate quantitative discipline based on precise observational data (see e.g. textbooks [4, 78]). Today, cosmology has its own "standard model" which is in a good agreement with most observational data. The basis of the model is a concept of the expanding Universe which, long ago, was very dense and so hot that the energy of thermal motion of elementary particles did not allow them to compose bound states. As a result, it were the particle interactions which determined all processes and, in the end, influenced the Universe development and the state of the world as we observe it today. The expanding Universe cooled down and particles were unified into bound states – first atomic nuclei from nucleons, then atoms from nuclei and electrons. Unstable particles decayed and the Universe arrived to its present appearence. As we will see below, presently the Universe expands with acceleration and is comprised mainly from unknown particles.

Even a dedicated book would be insufficient to describe all aspects of interrelations between cosmology and particle physics (the readers of *Physics Uspekhi* might be interested in reviews [79, 80]). Here, we will briefly consider three principal observational indications in favour of physics beyond SM, namely, the baryon asymmetry of the Universe, the dark matter and the accelerated expansion of the Universe (both the related notion of the dark energy and physical reasons for inflation).

3.1 Baryon asymmetry.

Quark-antiquark pairs had to be created intensively in the hot early Universe. The Universe then expanded and cooled down, quarks and antiquarks annihilated and the survived ones composed baryons (protons and neutrons). Notably, there are very few antibaryons in the present Universe, which means that at the early stages, there were more quarks than antiquarks. One can determine, by which amount: the number of quark-antiquark pairs was of the same order as the number of photons while the baryon-photon ratio may be determined from the analysis of the cosmic microwave background anisotropy and from studies of primordial nucleosynthesis. The ratio of the excess of quark number n_q over the antiquark number, $n_{\bar{q}}$, is of order

$$\frac{n_q - n_{\bar{q}}}{n_q + n_{\bar{q}}} \sim 10^{-10},$$

that is a single "unpaired" quark was present for each ten billion quark-antiquark pairs. It is hard to imagine that this tiny excess of matter over antimatter was present in the Universe from the very beginning; moreover, a number of quantitative cosmological models predict exact baryon symmetry of the very early Universe. Looks like the asymmetry appeared in course of the evolution of the Universe. For this to happen, the following Sakharov conditions [81] should be fulfilled:

1. baryon number nonconservation;
2. CP violation;
3. breaking of thermodynamical equilibrium.

Though the classical SM lagrangian conserves the baryon number, nonperturbative quantum corrections may break it, that is the condition 1 may be fulfilled in SM. The source of CP violation (condition 2) is also present in SM, it is the phase in the quark mixing matrix. Finally, in the course of the evolution of the Universe, the state

with the zero vacuum expectation value of the Higgs field (high temperature) has been replaced by the present state. It can be shown (see e.g. [82] and references therein), that the thermodynamic equilibrium was strongly broken at that moment, if it were a first-order phase transition. Therefore, in principle, all three conditions might be met in SM. However, it has been shown that the first-order electroweak phase transition in SM is possible only for the Higgs boson mass $M_H \lesssim 50$ GeV which was excluded from direct searches long ago. Also, the amount of CP violation in the CKM matrix is insufficient. We conclude that the observed baryon asymmetry of the Universe is an indication to the incompleteness of SM. A particular mechanism of generation of the baryon asymmetry is yet unknown (it should also explain the smallness of the asymmetry amount, $\sim 10^{-10}$).

3.2 Dark matter.

Study of dynamics of astrophysical objects (galaxies, galaxy clusters) and of the Universe as a whole allows one to determine the distribution of mass which may be subsequently compared to the distribution of the visible matter. Various independent observational data point to the estimate that the contribution of the visible matter (mostly baryons) to the energy density of the Universe is five times smaller than the contribution of invisible matter. We will first briefly discuss modern observational evidence for existence of the dark matter and then proceed to the discussion of implications of these observations to the particle physics.

1. Rotation curves of galaxies. Serious attention has been attracted to the question of the invisible matter after the analysis of rotation curves of galaxies (see e.g. [83]) (Fig. 7). For nearby galaxies it is possible to measure, by making use of the Doppler effect, the velocities of stars and gas clouds at different distances from the galaxy center, that is from the rotation axis. The Newton law of gravitation allows to estimate the distribution of mass as a function of the distance from the center; it was found that in the outer parts of galaxies, where luminous matter is practically absent, there is a significant mass density, so that the visible part of a galaxy is embedded into a much larger invisible massive halo. These measurements have been performed for many galaxies, in particular for our Milky Way.

2. Dynamics of galaxy clusters. In a similar way (though based on completely different observations), it is possible to determine the mass distribution in galaxy clusters. This provided for the historically first argument in favour of dark matter [86]. Modern observations have demonstrated that the main part of the baryonic matter sits not in the star systems, galaxies, but in hot gas clouds in the intergalactic space. This gas emits X rays, so the observations allow to reconstruct the distribution of the electron density and temperature. From the latter, by making use of the conditions of hydrostatic equilibrium, the mass distribution may be determined. Comparison with the distribution of the luminous matter (that is, mostly of the gas) points again to the existence of some hidden mass. A similar, though less precise, conclusion may be obtained from the analysis of velocities of galaxies inside a cluster.

3. Gravitational lensing. It may happen that a massive object (e.g. a galaxy cluster) is located between a distant source (e.g. a galaxy) and the observer. According to the general relativity, the light from the source is deflected by the massive object, so the latter may serve as a gravitational lens, which produces several distorted images of the source. A joint analysis of images of several sources allows one to model the mass distribution in the lens and to compare it with the distribution of visible matter (see e.g. [87]). The baryon distribution is reconstructed from X-ray observations of the



Figure 7: Some of the first indications to the existence of dark matter have been obtained from the analysis of the rotation curves of galaxies. Observational data on the rotation velocity as a function of the distance to the axis, given here for the galaxy NGC 3198 (dots), are not described by the curve which represents the expected velocity calculated from the distribution of luminous matter (lower line; data and calculation from [84]). At distances $\gtrsim$ 10 kpc, the luminous matter is practically absent (as one may see at the lower photograph taken from the digitalized Palomar sky atlas [85]), but the rotation velocities of gas clouds seen in the radio band are almost constant. This indicates that at periphery of the galaxy, there is a significant concentration of mass (the so-called halo).



Figure 8: The galaxy cluster Abell 1689. The background image of the cluster in the optical band was obtained by the Hubble Space Telescope (image from the archive [88]). The contours describe the model of mass distribution (full curves, Ref. [87]) based on the gravitational lensing and the distribution of luminous gas observed in X rays (dotted curves based on data from the Chandra X-ray telescope archive, Ref. [89]). Currently this mass model is one of the most precise ones.

luminous gas which contains about 90% of the cluster mass (Fig. 8). The full mass of the cluster calculated in this way far exceeds the mass of the baryons obtained from observations.

4. Colliding clusters of galaxies. One of the most beautiful observational proofs of the existence of dark matter is [90] the observation of colliding clusters of galaxies (Fig. 9). Contrary to the case of a usual cluster, Fig. 8, one does not need to calculate the mass in this case: comparison of the mass distribution and the gas distribution demonstrates that the main part of the mass of the clusters and that of luminous matter are located in different places. The reason for this dramatic difference, not seen in normal, noninteracting clusters, is related to the fact that the dark matter, constituting the dominant part of the mass, behave as a nearly collisionless gas. During the collision of clusters, the dark matter of one cluster, together with rare – and therefore also collisionless – galaxies kept by its gravitational potential, went through another one, while the gas clouds collided, stopped and were left behind.

These results, both by themselves and in combination with other results of quantitative cosmology (first of all those obtained from the analysis of the cosmic microwave background and the large-scale structure of the Universe, see e.g. [4]), point unequivocally to the existence of nonluminous matter. One should point out that the terms "dark", or "nonluminous", mean that the matter does not interact with the electromagnetic radiation and not only happens to be in a non-emitting state. Indeed, it should not also absorb electromagnetic waves, since otherwise the induced radiation would inevitably appear. The usual matter, that is baryons and electrons, may be put in this state only if packed in compact, very dense objects (neutron stars, brown



Figure 9: As in Fig. 8 but for colliding clusters 1E 0657–558 (the mass distribution model from [91], optical and X-ray images from archives [88, 89] correspondingly). Squares denote the positions of maxima of the mass distributions; diamonds denote the positions of maxima of the gas emission.

dwarfs etc.) which should be located in the halo of our Galaxy, as well as in other galaxies and in the intergalactic space within clusters. One may estimate the amount of these objects which is required to explain the observational results concerning the nonluminous matter. This amount appears to be so large that these compact objects should often pass between the observer and some distant sources, which should result in temporal distortion of the source image because of the gravitational lensing (the so-called microlensing effect). These events have been indeed observed, but at a very low rate which allows for a firm exclusion of this explanation for dark matter [92].

We are forced to say that, probably, the dark matter consists of new, yet unknown, particles, so that its explanation requires to extend SM. The dark-matter particles should be (almost) stable in order not to decay during the lifetime of the Universe ($\sim$14 billion years). These particles should also interact with the ordinary matter only very weakly to avoid direct experimental detection (direct searches for the dark matter, which should exist everywhere, in particular in laboratories, last already for decades). A number of theoretical models of the dark-matter origin predict the mass of the new particle between ~ 1 GeV and ~ 1 TeV and the cross section of interaction with ordinary particles of order of a typical weak-interaction cross section. Particles with these properties are called WIMPs (weakly interacting massive particles); they are absent in SM but exist in some of its extensions. One of the most popular candidates for the WIMP is the lightest superpartner (LSP) in supersymmetric extensions of SM with conserved R parity (see Sec. 4.2 below). The LSP cannot decay because the conservation of R parity requires that among decay products, at least one supersymmetric particle should be present, while all other supersymmetric particles are heavier by definition (in the same way, the electric-charge conservation provides for the electron stability and the baryon-number conservation provides for the stability of the proton). In a wide class of models the LSP is an electrically neutral particle (neutralino) which is considered a good candidate for a dark-matter particle. Note that there is a plethora of other scenarios in which the dark-matter particles have very different masses, from 10^{-5} eV (axion) to 10^{22} eV (superheavy dark matter). Also, in principle, the dark matter may consist of large composite particles (solitons).

3.3 Accelerated expansion of the Universe

In this section, we briefly discuss several technically interrelated problems which concern one of the least understandable, from the particle-physics point of view, part of the modern cosmology. They include:

1. observation of the accelerated expansion of the Universe ("dark energy");
2. weakness of the effect of the accelerated expansion as compared to typical scales of particle physics (the cosmological-constant problem);
3. indications to intense accelerated expansion of the Universe at one of the early stages of its evolution (inflation).

Let us start with the observational evidence in favour of (recent and present) accelerated expansion of the Universe.

1. The Hubble diagram. The first practical instrument of quantitative cosmology, the Hubble diagram plots distances to remote objects as a function of the cosmological redshift of their spectral lines. It was the way to discover the expansion of the Universe and to measure its rate, the Hubble constant. When methods to measure distances to the objects located really far away became available for astronomers, they found (see e.g. [93, 94]) deviations from a simple Hubble law which indicate that the expansion rate of the Universe changes with time, namely the expansion accelerates. The method of distance determination we are speaking about[2] is based on the study of type Ia supernovae and deserves a brief discussion (see also [95]).

A probable mechanism of the type Ia supernova explosion is the following. A white dwarf (a star at the latest stage of its evolution in which nuclear reactions have stopped) is rotating in a dense double system with a normal star. The matter from the normal star flows to the white dwarf and increases its mass. When the mass exceeds the so-called Chandrasekhar limit (the limit of stability of a white dwarf whose value depends, in practice, on the chemical composition of the star only), intense thermonuclear reactions start and the white dwarf explodes. It is interesting and useful to note that, therefore, in all cases the exploding stars have roughly the same mass and constitution (up to details of the chemical composition). As a consequence, all type Ia supernova explosions resemble each other not only qualitatively, but quantitatively as well: the energy release is roughly the same; the time dependence of the luminosity is similar. Even more amusing is the fact that even for rare outsiders (which differ from the most of supernovae either by the chemical composition or by some random circumstances), all curves representing the luminosity as a function of time are homothetic (Fig. 10), that is map one to another at a simultaneous scaling of both time and luminosity. It means that, upon measurement of a lightcurve of any type Ia supernova, one may determine its absolute luminosity with a good precision. Then, comparison with the observed magnitude allows to determine the distance to the object. In this way it is possible to construct the Hubble diagram (Fig. 11) which demonstrates statistically significant deviations from the law which corresponds to the uniform (or decelerated) expansion of the Universe.

2. Gravitational lensing. The method of the gravitational lensing, discussed above, allows not only for reconstruction of the mass distribution in the lensing cluster of galaxies, but also for determination of geometrical distances between sources, the lens and the observer. If redshifts of the source and the lens are known, one may compare

[2]The Nobel prise, 2011.



Figure 10: Temporal dependence of the absolute magnitude of type Ia supernovae. *Above:* lightcurves of 68% of supernovae are contained within the shaded band, however there are very rare outsiders (for example, lightcurves of an unusually bright supernova SN1991T (squares) and an unusually weak supernova SN1986G (triangles) are presented); light curves and the band are taken from [96]. *Below:* the same curves but scaled simultaneously in the horizonthal (time) and vertical (luminosity) axes according to the rules described in [97]. Introduction of this correction shifts all "exclusive" curves to the band. Therefore, to know the absolute value of the luminosity, it is sufficient to measure the shape of the light curve.



Figure 11: The Hubble diagram, presenting the dependence of the distance from the redshift z of spectral lines of distant galaxies, obtained from observations of type Ia supernovae. Gray lines correspond to data on individual supernovae with experimental error bars [98]. The uniform expansion of the Universe corresponds to the lower (dotted) line, the accelerated expansion – to the upper (full) line.

them with the derived distances and find [99] deviations from the Hubble law with a high precision.

3. Flatness of the Universe and the energy balance. A number of measurements point to the spatial flatness of the Universe, that is to the fact that its three-dimensional curvature is zero. The main argument here is based on the analysis of the cosmic microwave background anisotropy [100]. In the past, the Universe was denser and hotter than now. Various particles (photons in particular) were in thermodynamical equilibrium, so that the distribution of photons over energies was Planckian, corresponding to the temperature of the surrounding plasma. The Universe cooled down while expanding and at some moment, electrons and protons started to join into hydrogen atoms. Compared to plasma, the gas of neutral atoms is practically transparent for radiation; since then, photons born in the primordial plasma propagate almost freely. We see them as the cosmic microwave background (CMB) now. At the moment when the Universe became transparent, the size of the causally connected region (that is the region which a light signal had time to cross since the Big Bang), called a horizon, was only ~ 300 kpc. This quantity may be related to a typical scale of the CMB angular anisotropy; the present Universe is much older and we see at the same moment many regions which had not been causally connected in the early Universe. This angular scale has been directly measured from the CMB anisotropy. The theoretical relation between this scale and the size of the horizon at the moment when the Universe became transparent is very sensitive to the value of the spatial curvature; the analysis of the data from WMAP satellite points to a flat Universe with a very high accuracy.

Other methods exist to test the flatness of the Universe. One of the most beautiful among them is the geometric Alcock-Paczinski criterion. If it is known that an object has a purely spherical shape, one may try to measure its dimensions along the line of sight and in the transverse direction. Taking into account distortions related to the expansion of the Universe, one may compare the two sizes and constrain the cosmological parameters, first of all, deviations from flatness. Clearly, it is not an easy task to find an object whose longuitudinal and transverse dimensions are certainly equal; however, one may measure characteristic dimensions of some astrophysical structures which, averaged over large samples, should be isotropical. The most precise measurement of this kind [101] uses double galaxies whose orbits are randomly oriented in space while the orbital motion is described by the Newton dynamics.

From the general-relativity point of view, the flat Universe represents a rather specific solution which is characterized by a particular total energy density (the so-called critical density, $\rho_c \sim 5 \times 10^{-6}\ \frac{\mathrm{GeV}}{\mathrm{cm}^3}$). At the same time, the estimates of the energy density related to matter contribute $\sim 0.25\rho_c$, that is the remaining three fourths of the energy density of the Universe are due to something else. This contribution, whose primary difference form the matter contribution is in the absence of clustering (that is of concentration in stars, galaxies, clusters etc.), carries a not fully successful name of "dark energy".

The question about the nature of the dark energy is presently open. The technically most simple explanation is that the accelerated expansion of the Universe results from a nonzero vacuum energy (in general relativity, the reference point on the energy axis is relevant!), that is the so-called cosmological constant. From the particle-physics point of view, the dark-energy problem is, in this case, twofold. In the absence of special cancellations, the vacuum energy density should be of order of the characteristic scale of relevant interactions Λ, that is

$$\rho \sim \frac{\Lambda^4}{c^3\hbar^3}.$$

The observed value of ρ corresponds to $\Lambda \sim 10^{-3}$ eV, while characteristic scales of the strong ($\Lambda_{\rm QCD} \sim 10^8$ eV) and electroweak ($v \sim 10^{11}$ eV) interactions are many orders of magnitude higher. One side of the problem (known for a long time as "the cosmological-constant problem") is to explain how the contributions of all these interactions to the vacuum energy cancel. In principle, some symmetry may be responsible for this cancellation: for instance, the energy of a supersymmetric vacuum in field theory is always zero. Unfortunately, the supersymmetry, even if it has some relation to the real world, should (as discussed in Sec. 4), be broken at the scale not smaller than $\sim v$, and the contributions to the vacuum energy should have then the same order. On the other hand, the observed accelerated expansion of the Universe tells us that the cancellation is not complete and hence there is a new energetic scale in the Universe $\sim 10^{-3}$ eV. The explanation of this scale is a task which cannot be completed within the frameworks of SM, where all parameters of the dimension of energy are orders of magnitude higher. If this scale is given by a mass of some particle, the properties of the latter should be very exotic in order both to solve the problem of the accelerated expansion of the Universe and not to be found experimentally. For instance, one of suggested explanations [102] introduces a scalar particle whose effective mass depends on the density of medium (this particle is called "chameleon"). By itself, the dependence of the effective mass on the properties of medium is well-known (for instance, the dispersion relation of photon in plasma is modified in such a way that it gets a nonzero effective mass). In our case, due to interaction with the external gravitational field, the chameleon has a short-distance potential in relatively dense medium (e.g. at the Earth), which prohibits its laboratory detection, but at large scales of the (almost empty) Universe the effect of this particle becomes important. One should also note that a solution to the problem of the accelerated expansion of the Universe might have nothing to do with particle physics at all and be entirely based on peculiar properties of the gravitational interaction (for instance, on deviations from the general relativity at large distances).

However, the problem of the accelerated expansion of the Universe is not exhausted by the analysis of the modern state. There are serious indications that, at an early stage of its evolution, the Universe experienced a period of intense exponential expansion, called inflation (see e.g. [78, 103]. Though the inflation theory is currently not a part of the standard cosmological model (it awaits more precise experimental tests), it solves a number of problems of the standard cosmology and, presently, does not have an elaborated alternative. Let us list briefly some problems which are solved by the inflationary model.

1. As it has been already pointed out, various parts of the presently observed Universe were causally disconnected from each other in the past, if one extrapolates the present expansion of the Universe backwards in time. Information between the regions which are now observed in different directions could not be transmitted, for instance, at the moment when the Universe became transparent for CMB. At the same time, the CMB is isotropic up to a high level of accuracy (relative variations of its temperature does not exceed 10^{-4}), the fact that indicates to the causal connection between all currently observed regions.

2. Zero curvature of the Universe, from the theoretical point of view, is not singled out by any condition: the Universe should be flat from the very beginning, nobody knows why.

3. The modern Universe is not precisely homogeneous – the matter is distributed inhomogeneously, being concentrated in galaxies, clusters and superclusters of

galaxies; a weak anisotropy is observed also in CMB. Most probably, these structures were developed from tiny primordial inhomogeneities, whose existence should be assumed as the initial condition.

These and some other arguments point to the fact that the initial conditions for the theory of a hot expanding Universe had to be very specific. A simultaneous solution to all these problems is provided by the inflationary model which is based on the assumption about an exponential expansion of the Universe which happened before the hot stage. From the theory point of view, this situation is fully analogous to the present accelerated expansion, but the energy density, which determines the acceleration rate, was much higher. It may be related to the presence of a new, absent in SM, scalar field, the inflaton. If it has a relatively flat (that is, weakly depending from the field value) potential and the value itself slowly changes with time, then the energy density of the inflaton provides for the required exponential expansion. For a particle physicist, at least two questions arise: first, what is the nature of the inflaton, and second, what was the reason for the inflation to stop and not to continue until now.

To summarize, we note that a large number of observations related to the structure and evolution of the Universe cannot be explained if the particle physics is described by SM only: one needs to introduce new particles and interactions. Jointly with the observation of neutrino oscillations, these facts constitute the experimental basis for the confidence in incompleteness of SM. At the same time, presently none of these experimental results point to a specific model of new physics, so one is guided also by purely theoretical arguments when constructing hypothetical models.

4 Aesthetic difficulties: the origin of parameters.

4.1 Electroweak interaction and the Higgs boson.

Results of high-precision measurements of electroweak-theory parameters, in particular at the LEP accelerator, confirm the predictions of SM, based on the Higgs mechanism. At the same time, the only SM particle which has not been discovered experimentally, is the Higgs boson. Its mass is a free parameter of the model and is not directly related to any of measurable parameters, so the lack of signs of the Higgs boson in data may be simply explained by its mass: the energies and luminosities of available accelerators might be insufficient to create this particle with a significant probability.

At the same time, purely theoretical concerns suggest that the Higgs boson should not be too heavy. It is related to the fact that, without the account of the Higgs scalar, the scattering amplitudes of massive W bosons grow as E^2 with energy E. As a result, at energies somewhat higher than the W mass, the perturbation theory fails and all model predictions start to depend on unknown higher-order contributions; the theory finds itself in a strong-coupling regime and loses predictivity. The contribution from the Higgs boson, however, cancels the part of the amplitude which grows with energy, so only the constant term remains, $\sim g^2 M_H^2/(4M_W^2)$, where M_H and M_W are masses of the Higgs and W bosons, respectively, and g is the $SU(2)_{\rm L}$ gauge coupling constant. Therefore, to keep calculability, M_H should not be too large; a quite reliable limit is $M_H \lesssim 800$ GeV. Even more restrictive limits come from the radiative corrections to the potential of the Higgs boson itself. In the leading order of perturbation theory, the self-interaction constant of the Higgs boson has a pole at the energy scale

$$Q \sim v \exp\left[\frac{4\pi^2 v^2}{3M_H^2}\right].$$



Figure 12: The Higgs boson mass expected from indirect data and constrained from direct searches (see text). The left panel shows all experimental limits; the right one is a zoom of the most interesting region, $M_H < 200$ GeV.

This means that at energies $\Lambda \leq Q$, the contributions of new particles or interactions should change the coupling behaviour to avoid divergence. The requirement $\Lambda \geq 1$ TeV results in the limit $M_H \lesssim 550$ GeV. Note that this means that the SM Higgs boson should be discovered at LHC.

Maybe even more interesting situation is related to the experimental data on the search of the Higgs particle. The latter may reveal itself not only directly, being produced at colliders, but also indirectly, through the influence of the virtual Higgs bosons on numerous observables. Though this influence is not large, a numebr of electroweak observables are known with a very high precision, so that their joint analysis may constrain the mass of yet undiscovered Higgs particle. Let us look at Fig. 12, which is based on the analysis of indirect experimental data[3] as of September 2011. The horizontal axis gives the possible Higgs-boson mass; the shaded regions of M_H are excluded, as of December 2011, from direct experimental search of the Higgs boson at colliders at the 95% confidence level (the light band $M_H < 114$ GeV – LEP [104], the light bands 114 GeV$< M_H <$ 115.5 GeV and 127 GeV$< M_H <$ 600 GeV – LHC [105, 106], the dark bands 100 GeV$< M_H <$ 109 GeV and 156 GeV$< M_H <$177 GeV – Tevatron [107]). The curve [108] demonstrates how well a given value of M_H agrees with a combination of all *other* than the direct-search experiments (the lower $\Delta\chi^2$, the better agreement; the curve width represents the uncertainty in theoretical predictions). One can see that the most preferable value of M_H is already experimentally excluded! Clearly, this does not mean a catastrophe because a narrow range of slightly less preferable values are allowed, but it motivates theoretical physicists to think about possible alternative explanations of the electroweak symmetry breaking [109]. One should note that it is rather difficult to discover a light, 115 GeV$< M_H <$ 127 GeV, Higgs boson at LHC: unlike for a heavy one, several years of work might be required.

The lack of the Higgs boson with the expected mass and the prospect of further restriction of the allowed mass region at LHC are important, but far not principal arguments in favour of alternative theoretical models of the electroweak symmetry breaking, whose history goes back for decades. The point is that the Higgs boson is the only SM scalar particle (all others are either fermions or vectors). A scalar particle brings to a theory a number of unfavoured properties some of which we have just mentioned above, while others will be discussed below. That is why alternative mechanisms of the electroweak symmetry breaking use, as a rule, only fermionic and

[3]See also the regularly updated webpage at http://gfitter.desy.de/GSM .

vector fields.

A class of hypothetical models in which the vacuum expectation value of the Higgs particle is replaced by the vacuum expectation value of a two-fermion operator with the same quantum numbers are called technicolor models (see e.g. [110]). The replacement of the scalar by a fermion condensate looks quite natural if one recalls that in the historically first, and surely realized in Nature, example of the Higgs mechanism, the Ginzburg–Landau superconductor, the condensate of the Cooper pairs of electrons plays the role of the Higgs boson.

The base for the construction of technicolor models is provided by the analogy to QCD. Indeed, an unbroken nonabelian gauge symmetry, similar to $SU(3)_{\rm C}$, may result in confinement of fermions and to formation of bound states (in QCD these are hadrons, bound states of quarks). In fact, in QCD a nonzero vacuum expectation value of the quark condensate also appears, but its value, of order $\Lambda_{\rm QCD}$ ∼200 MeV, is much less than the required electroweak symmetry breaking scale ($v \approx 246$ GeV). Therefore one postulates that there exists another gauge interaction, in a way resembling QCD, but with a characteristic scale of order v. The corresponding gauge group $G_{\rm TC}$ is called a technicolor group. The bound states, technihadrons, are composed from the fundamental fermions, techniquarks T, which feel this interaction. The techniquarks carry the same quantum numbers as quarks, except instead of $SU(3)_{\rm C}$, they transform as a fundamental representation of $G_{\rm TC}$. Then, the vacuum expectation value $\langle \bar{T}T \rangle$ breaks $SU(2)_{\rm L} \times U(1)_{\rm Y} \to U(1)_{\rm EM}$ in such a way that the correct relation between masses of the W and Z bosons is fulfilled automatically. A practical implementation of this beautiful idea faces, however, a number of difficulties which result in a complification of the model. First, the role of the Higgs boson in SM is not only to break the electroweak symmetry: its vacuum expectation value also gives masses to all charged fermions. Attempts to explain the origin of fermion masses in technicolor models result in significant complication of the model and, in many cases, in contradiction with experimental constraints on the flavour-changing processes. Second, many parameters of the electroweak theory are known with very high precision (and agree with the usual Higgs breaking), while even a minor deviation from the standard mechanism destroys this well-tuned picture. To construct an elegant and viable technicolor model is a task for future which will become relevant if the Higgs scalar will not be found at LHC.

In another class of models (suggested in [111] and further developed in numerous works which are reviewed, e.g., in [109]), the Higgs scalar appears as a component of a vector field. Since the vacuum expectation value of a vector component breaks Lorentz invariance, this mechanism works exclusively in models with extra space dimensions. For instance, from the four-dimensional point of view, the fifth component of a five-dimensional gauge field behaves as a scalar, and giving a vacuum expectation value to it breaks only five-dimensional Lorentz invariance while keeping intact the observed four-dimensional one. Symmetries of the five-dimensional model, projected onto the four-dimensional world, protect the effective theory from unwanted features related to the existence of a fundamental scalar particle. These models also have a number of phenomenological problems which can be solved at a price of significant complication of a theory.

The so-called higgsless models [112] (see also [109]) are rather close to these multi-dimensional models, though differ from them in some principal points. The higgsless models are based on the analogy between the mass and the fifth component of momentum in extra dimensions: both appear in four-dimensional effective equations of motion similarly. In the higgsless models, the nonzero momentum appears due to imposing some particular boundary conditions in a compact fifth dimension. In the end,

these boundary conditions are responsible for breaking of the electroweak symmetry. Unlike in five-dimensional models, where the Higgs particle is a component of a vector field, the physical spectrum of the effective theory in higgsless models does not contain the corresponding degree of freedom. These models have some phenomenological difficulties (related e.g. to precise electroweak measurements). Another shortcoming of this class of models is considerable arbitraryness in the choice of the boundary conditions, which are not derived from the model but are crucial for the electroweak breaking.

Finally, we note that a composite Higgs boson may be even more complex than just a fermion condensate: it may be a bound state which includes strongly coupled gauge fields. Description of these bound states requires a quantitative understanding of nonperturbative gauge dynamics. Taking into account the analogy between strongly coupled four-dimensional theories and weakly coupled five-dimensional ones (which will be discussed in Sec. 5.3, these models may even happen to be equivalent to multidimensional models described above.

4.2 The gauge hierarchy.

Each of the main interactions of particles has its own characteristic energy scale. For the strong interaction it is $\Lambda_{\rm QCD} \sim 200$ MeV, the scale at which the QCD running coupling becomes strong; this scale determines masses of hadrons made of light quarks. The scale of the electroweak theory is determined by the vacuum expectation value of the Higgs boson, $v \approx 246$ GeV, which determines, through the corresponding coupling constants, the masses of the W and Z bosons and of SM matter fields. For gravity, the characteristic scale is the Planck scale $M_{\rm Pl} \sim 10^{19}$ GeV, determined by the Newton constant of the classical gravitational interactions.

These three scales are related to known forces. Extensions of SM give motivation to some other interactions and, consequently, to other scales. First of all it is $M_{\rm GUT} \sim 10^{16}$ GeV, the scale of the suggested Grand Unification of interactions. In several models explaining neutrino masses there exists a scale M_ν; sometimes the scale $M_{\rm PQ}$, related to the CP invariance of the strong interaction, is also introduced. Values of these two scales are model dependent but roughly $M_{\rm PQ} \sim M_\nu \sim 10^{14}$ GeV.

The gauge hierarchy problem (see also [2, 79, 113]) consists in the disproportionality of these scales:

$$(\Lambda_{\rm QCD}, v) \ll (M_{\rm Pl}, M_{\rm GUT}, M_{\rm PQ}, M_\nu)$$

and in a range of related questions which may be divided into three groups.

1. The origin of the hierarchy: why the scales of the strong and electroweak interactions are smaller than others by many orders of magnitude? That is, why, for instance, all SM particles are practically massless at the gravity scales? It is possible, in the frameworks of the Grand-Unification hypothesis, to get a reasonable explanation of the relation $\Lambda_{\rm QCD} \ll M_{\rm GUT}$. It is based on the logarithmic renormalization-group dependence of the gauge coupling constant from energy E. In the leading approximation, this dependence for the strong-interaction coupling α_3 reads

$$\alpha_3(E) = \frac{\alpha_{\rm GUT}}{1 + \beta_3 \alpha_{\rm GUT} \ln(E/M_{\rm GUT})},$$

where β_3 is a positive coefficient which depends on the set of strongly interacting matter fields (in SM, $\beta_3 = 11/(12\pi)$), while $\alpha_{\rm GUT} \sim 1/30$ is the value of the coupling constant of a unified gauge theory at the energy scale $\sim M_{\rm GUT}$. The scale $\Lambda_{\rm QCD}$, where α_3



Figure 13: Hierarchy of scales of gauge interactions.

becomes large, may be determined in this approximation as

$$\Lambda_{\rm QCD} = M_{\rm GUT} \exp\left(-\frac{1}{\beta_3 \alpha_{\rm GUT}}\right)$$

and the exponent provides for the required hierarchy. However, a similar analysis is not succesful for the electroweak interaction, whose coupling constants are small at the scale v. The latter is unrelated to any dynamical scale and is introduced in the theory as a free parameter.

2. The stability of the hierarchy. In the standard mechanism of the electroweak breaking, the characteristic scale $v = M_H/\sqrt{2\lambda}$, where λ is the self-interaction constant of the Higgs boson. Together with M_H, the scale v gets, in SM, quadratically divergent radiative corrections,

$$\delta v^2 \sim \delta M_H^2 = f(g)\Lambda_{\rm UV}^2,$$

where $f(g)$ is a symbolic notation for some known combination of the coupling constants (in SM, $f(g) \approx 0.1$), and $\Lambda_{\rm UV}$ is the ultraviolet cutoff which may be interpreted as an energy scale above which SM cannot give a good approximation to reality. This scale may be related to one of the scales $M_{\rm Pl}, M_{\rm GUT}$ etc. discussed above; in the assumption of the absence of the "new physics", that is of particle interactions other than those already discovered (SM and gravity), one should take $\Lambda_{\rm UV} \sim M_{\rm Pl}$. Therefore, since $v^2 = v_0^2 - \delta v^2$, where v_0 is the parameter of the tree-level lagrangian, the hierarchy $v^2 \ll M_{\rm Pl}^2$ appears as a result of cancellation between two huge contributions, v_0^2 and δv^2. Each of them is of order $f(g)M_{\rm Pl}^2 \sim 10^{33} v^2$, that is the cancellation has to be precise up to 10^{-33} in every order of the perturbation theory. This *fine tuning* of parameters of the model, though technically possible, does not look natural. One may revert this logic and say that to avoid fine tuning in SM one should have

$$f(g)\Lambda_{\rm UV}^2 \sim v^2 \quad \Rightarrow \quad \Lambda_{\rm UV} \sim {\rm TeV}. \tag{3}$$

The relation (3) gives a base for the optimism of researchers who expect the discovery of not only the Higgs boson but also some new physics beyond SM from the LHC.

3. The gauge desert. The third aspect of the same problem is related to the presumed absence of particles with masses (and of interactions with scales) between "small" $(\Lambda_{\rm QCD}, v)$ and "large" $(M_\nu, M_{\rm GUT}, M_{\rm Pl})$ energetic scales, see Fig. 13. All known particles are settled in a relatively narrow region of masses $\lesssim v$, beyond which, for many orders of magnitude, lays the so-called *gauge desert*. Clearly, one may suppose that the

heavier particles simply cannot be discoverd due to insufficient energy of accelerators, but this suggestion is not that easy to accomodate within the standard approach. Indeed, new relatively light ($\sim v$) particles which carry the SM quantum numbers are constrained by the electroweak precision measurements. Also, the latest Tevatron and first LHC results on the direct search of new quarks strongly constrain the range of their allowed masses (see [114] and references therein). In particular, for the fourth generation of matter fields similar to the known three, the mass of its up-type quark should exceed 338 GeV, while that of a down-type quark should exceed 311 GeV. The mass of the corresponding charged lepton cannot be lower than 101 GeV [6]. The mass of the fourth-generation standard neutrino should exceed one half of the Z-boson mass, as it has been already discussed above. At the same time, these values of masses of the fourth-generation charged fermions cannot have the same origin as those for the first three generation, because to generate masses much larger than v, Yukawa constants much larger than one are required. Since the methods to calculate nonperturbative corrections to masses are yet unknown, for this case one cannot be sure that these masses can be obtained at all in a usual way. Moreover, SM fermion masses exceeding the electroweak breaking scale are forbidden by the $SU(2)_L \times U(1)_Y$ gauge symmetry: a mechanism generating these masses would also break the electroweak symmetry at a scale $> v$. Addition of matter fields which do not constitute full generations may be considered as an essential extension of SM. Finally, addition of new matter affects the energy dependence of the gauge coupling constants and spoils their perturbative unification (unless one adds either full generations or other very special sets of particles of roughly the same mass which constitute full multiplets of a unified gauge group). We see that attempts "to inhabit the gauge desert" inevitably result in significant steps beyond SM while the desert itself does not look natural.

Attempts to solve the gauge hierarchy problem may be also divided into several large groups.

1. The most radical approach, rather popular in recent years, is to assume that the high-energy scales are simply absent in Nature. For a theoretical physicist, the most easy scales to refuse are M_ν and $M_{\rm PQ}$, because they do not appear in all models explaining neutrino masses and CP conservation in strong interactions, respectively. $M_{\rm GUT}$ is a bit more difficult: the Grand Unification of interactions gets support not only from aesthetic expectations (electricity and magnetism unified to electrodynamics, electrodynamics and weak interactions unified to the electroweak theory, etc.) and the arguments related to the electric charge quantization (see e.g. [3]), but also from the analysis of the renormalization-group running of the three SM gauge coupling constants which get approximately the same value at the scale $M_{\rm GUT}$ (see e.g. [2, 3]). It is worth noting that at the plot (see Fig. 14) of $\alpha_{1,2,3}$ as functions of energy in SM, the three lines do not intersect at a strictly one point, however, for the case of evolution for many orders of magnitude in the energy scale, already an approximate unification is a surprise. To make three lines intersect at one point precisely, one needs a free parameter, which may be introduced in the theory with some new particles, e.g. with masses $\sim$TeV (this happens in particular in models with low-energy supersymmetry, see below). Therefore, the most amusing is not the precise unification of couplings in an extended theory with additional parameters but the approximate unification already in SM. It is not that easy to keep this miraculous property and at the same time to lower the $M_{\rm GUT}$ scale in order to avoid the hierarchy $v \ll M_{\rm GUT}$. Indeed, the addition of new particles which affect the renormalization-group evolution either spoils the unification or, to the leading order, does not change the $M_{\rm GUT}$ scale (note that in SM, the unification occurs in the perturbative regime and higher corrections does not change the picture significantly).



Figure 14: The energy-scale dependence of coupling constants of SM gauge interactions $U(1)_{\rm Y}$ (the full line), $SU(2)_{\rm W}$ (the dashed line) and $SU(3)_{\rm C}$ (the dotted line) in the leading order.

The only possibility is to give up the perturbativity (the so-called "strong unification" [115, 116]). In this latter approach, the addition of a large number of new fields in full multiplets of a certain unified gauge group results in increasing of the coupling constants at high energies; QCD stops to be asymptotically free at energies higher than the masses of the new particles. In the leading order, all three coupling constants have, in this case, poles at high energies; the unification of SM couplings guarantees that the three poles coincide and are located at $M_{\rm GUT}$. However, this leading-order approximation has nothing to do with the real behaviour of constants in the strong-coupling regime, so the theory may generate a new scale M_s at which $\alpha_{1,2,3}$ become strong, this scale being an ultraviolet analog of $\Lambda_{\rm QCD}$. For a sufficiently large number of additional matter fields, M_s may be sufficiently close to the electroweak scale v: in certain cases, it might be that $M_s \ll M_{\rm GUT}$ (a nonperturbative fixed point). In this scenario, low-energy observable values of the coupling constants appear as infrared fixed points and do not depend on unknown details of the strong dynamics. Note that the Grand-unified theory may have degrees of freedom very different from SM in this case.

In the recent decade, the models became quite popular in which the hierarchy problem is solved by giving up the large parameter $M_{\rm Pl}$. This parameter is related to the gravitational law and any attempt to change the parameter requires a change in the Newtonian gravity. This may be achieved, for instance, if the number of space dimensions exceeds three but, for some reason, the extra dimensions remain unseen (see e.g. a review [117]). Indeed, assume that the extra dimensions are compact and have a characteristic size $\sim R$, where R is sufficiently small. Then it is easy to obtain the relation

$$M_{\rm Pl}^2 \sim R^\delta M_{{\rm Pl},4+\delta}^{2+\delta}, \tag{4}$$

where δ is the number of extra space dimensions, $M_{{\rm Pl},4+\delta}$ is the fundamental parameter of the $(4+\delta)$-dimensional theory of gravity, while $M_{\rm Pl}$ now is the effective four-dimensional Planck mass. Already in the beginning of the past century, in works by Kalutza [118], subsequently developed by Klein [119], possible existence of these extra

dimensions, unobservable because of small R, has been discussed. This approach assumed that $R \sim 1/M_{\rm Pl}$ (and therefore $M_{\rm Pl} \sim M_{{\rm Pl},4+\delta}$) and has become well-known and popular in the second part of the 20th century in context of various models of string theory, which however did not result in succesful phenomenological applications by now. We will discuss, in a little more detail, another approach which allows to make R larger without problems with phenomenology. It is based on the idea of localization of observed particles and interactions in a 4-dimensional manifold of a $(4+\delta)$-dimensional spacetime [120, 121, 122].

From the field-theoretical point of view, the localization of a $(4+\delta)$-dimensional particle means that the field describing this particle satisfies an equation of motion with variables related to the observed four dimensions (call them $x_\mu,\ \mu = 0,1,2,3$) separated from those related to δ extra dimensions ($z_A,\ A = 1,\dots\delta$) and the solution for the z-dependent part is nonzero only in a vicinity (of the size $\sim \Delta$) of a given point in the δ-dimensional space (without loss of generality, one may consider the point $z = 0$), while the x-dependent part satisfies the usual four-dimensional equations of motion for this field. As a result, the particles described by the field move along the four-dimensional hypersurface corresponding to our world and do not move away from it to the extra dimensions for distances exceeding Δ. This may happen if the particles are kept on the four-dimensional hypersurface by a force from some extended object which coincides with the hypersurface. This solitonlike object is often called brane, hence the expression "braneworld". The readers of *Physics Uspekhi* may find a more detailed description of this mechanism in [117].

Based on the topological properties of the brane, localisation of light (massless in the first approximation) scalars and fermions in four dimensions[4] implies that many direct experimental bounds on the size of extra dimensions in a Kalutza-Klein-like model restrict now the region Δ accessible for the observed particles, instead of the size R of the extra dimension. In [124], it has been suggested to use this possibility, for $R \gg \Delta$, to remove, according to Eq. (4), a large fundamental scale $M_{\rm Pl}$ and the corresponding hierarchy. It has been pointed out that in this class of models, R is bound from above mostly by nonobservation of deviations from the Newtonian gravity at short distances; experiments now exclude the deviations at the scales of order 50 μm only [125] (it was $\sim$ 1 mm at the moment when the model was suggested). This allows, according to Eq. (4), to have $M_{{\rm Pl},4+\delta} \sim$TeV, that is almost of the same order as v. Models of this class are well studied from the phenomenological point of view but have two essential theoretical drawbacks. The first one is related to the apparent absence of a reliable mechanism of localization of *gauge* fields in four dimensions. The only known field-theoretical mechanism for that [126] is based on some assumptions about the behaviour of a multidimensional gauge theory in the strong-coupling regime. Though these assumptions look realistic, they currently cannot be considered as well-justified. The second difficulty is aesthetic and is related to the appearence of a new dimensionful parameter R: the hierarchy $v \ll M_{\rm Pl}$ happens to be simply reformulated in terms of a new unexplained hierarchy $1/R \ll M_{{\rm Pl},4+\delta}$.

To a large extent, these difficulties are overcome in somewhat more complicated models, in which the spacetime cannot be presented as a direct product of our four-dimensional Minkowski space and compactified extra dimensions [127, 128, 129]. The principal difference of this approach from the one discussed above is that the gravitational field of hte brane in extra dimensions is not neglected. For $\delta = 1$ and in the limit

[4]Note that recently, a fully analogous mechanism of localisation in one- or two-dimensional manifolds has been tested *experimentally* for a number of solid-state systems (the quatum Hall effect, topological superconductors and topological insulators, graphene), see e.g. [123].

of a thin brane, one obtains the usual five-dimensional general-relativity equations. These equations have, in particular, solutions with four-dimensional Poincare invariance. The metrics in these solutions is exponential in the extra-dimensional coordinate (the so-called anti-de-Sitter space),

$$ds^2 = \exp(-2k|z|)dx^2 - dz^2 \tag{5}$$

where ds^2 and dx^2 are the squares of the five-dimensional and usual four-dimensional (Minkowski) intervals, respectively. For a finite size z_c of the fifth dimension, the relation between the fundamental scales is now

$$M_{\rm Pl} \sim \exp(kz_c) M_{\rm Pl,5}.$$

If fundamental dimensionful parameters of the five-dimensional gravity satisfy $M_{\rm Pl,5} \sim k \sim v$, one may [129] explain the hierarchy $v/M_{\rm Pl}$ for $z_c \approx 37/k$, that is instead of the fine tuning with the precision of 10^{-16}, one now needs to tune the parameters up to ~ 0.1. It is interesting that in models of this kind with two or more extra dimensions, it is possible [130] to localize gauge fields on the brane in the weak-coupling regime, contrary to the case of the factorizable geometry.

2. A completely different approach to the problem of stabilization of the gauge hierarchy is to add new fields which cancel quadratic divergencies in expressions for the running SM parameters. The best-known realization of this approach is based on supersymmetry (see e.g. reviews [131, 132, 133]), which provides for the cancellation of divergencies due to opposite signs of ferminonic and bosonic loops in Feynman diagrams.

The requirement of supersymmetry is very restrictive for the mass spectrum of particles described by the theory. Namely, together with the observed particles, their superpartners, that is particles with the same masses and different spins, should be present. The absence of scalar particles with masses of leptons and quarks and of fermions with masses of gauge bosons means that unbroken supersymmetry does not exist in Nature. It has been shown, however, that it is possible to break supersymmetry while keeping the cancellation of quadratic divergencies. This breaking is called "soft" and naturally results in massive superpartners.

In the minimal supersymmetric extension of SM (MSSM; see e.g. [133]), each of the SM fields has a superpartner with a different spin: the Higgs boson corresponds to a fermion, higgsino; matter-field fermions correspond to scalar squarks and sleptons; gauge bosons correspond to fermions which transform in the adjoint representation of the gauge group and are called gauginos (in particular, gluino for $SU(3)_C$, wino and zino for the W and Z bosons, bino for the hypercharge $U(1)_Y$ and photino for the electromagnetic gauge group $U(1)_{\rm EM}$). For the theory to be selfconsistent (absence of anomalies related to the higgsino loops), and also to generate fermion masses in a supersymmetric way, the second Higgs doublet is introduced, which is absent in SM. The cancellation of quadratic divergencies may be easily seen in Feynman diagrams: in the leading order, closed fermion loops have the overall minus sign and cancel the contributions from loops of their superpartner bosons. This cancellation is precise as long as the masses of particles and their superpartners are equal; otherwise the contributions differ by an amount proportional to the difference between squared masses of superpartners, Δm^2. The condition of stability of the gauge hierarchy then requires that $\frac{g^2}{16\pi^2}\Delta m^2 \lesssim v^2$, where g is the coupling constant in the vertex of the corresponding loop (the maximal, $g \sim 1$, coupling constant is that of the top quark). We arrive to an important conclusion which motivates in part the current interest to phenomenological supersymmetry: if the problem of stabilization of the gauge hierarchy is solved by

supersymmetry, then the superpartner masses cannot exceed a few TeV, which means that they might be experimentally found in the nearest future.

The MSSM lagrangian, in the limit of unbroken supersymmetry, satisfies all symmetry requirements of SM, including the conservation of the lepton and baryon numbers. At the same time, the SM gauge symmetries do not forbid, for this set of fields, certain interaction terms which violate the lepton and baryon numbers. The coefficients at these terms should be very small in order to satisfy experimental constraints, for instance, those related to the proton lifetime. It is usually assumed that these terms are forbidden by an additional global symmetry $U(1)_R$. When supersymmetry is broken, this $U(1)_R$ breaks down to a discreet Z_2 symmetry called R parity. With respect to the R parity, all SM particles carry charges $+1$ while all their superpartners carry charges -1. The R-parity conservation leads to the stability of the lightest superpartner (see Sec. 3.2).

The soft supersymmetry-breaking terms are introduced in the MSSM lagrangian explicitly. They include usual mass terms for gaugino and scalars as well as trilinear interactions of the scalar fields. In addition to the SM parameters, about 100 independent real parameters are therefore introduced. In general, these new couplings with arbitrary parameters may result in nontrivial flavour physics. The absence of flavour-changing neutral currents and of processes with nonconservation of leptonic quantum numbers, as well as limits from the CP violation, narrow the allowed region of the parameter space significantly.

One may note the following characteristic features of the phenomenological supersymmetry.

(1). The coupling-constant unification at a high energy scale becomes more precise as compared to SM, if superpartners have masses $\sim v$ as required for the stability of the gauge hierarchy.

(2). In the same regime, the gauge desert between $\sim 10^3$ GeV and $\sim 10^{16}$ GeV is still present.

(3). In MSSM, there is a rather restrictive bound on the mass of the lightest Higgs boson. In the leading approximation of perturbation theory, it is $M_H < M_Z$. The account of loop corrections allow to relax it slightly, but in most realistic models $M_H < 150$ GeV is predicted. The absence of a light Higgs boson discussed in Sec. 4.1 is a much more serious problem for supersymmetric theories than for SM.

(4). The phenomenological model described above explains the stability of the gauge hierarchy but not its origin. The small parameter v/M, where $M = M_{\rm GUT}$ or $M = M_{\rm Pl}$, does not require tuning in every order of perturbtion theory but should be introduced in the model by hand, that is cannot be derived, nor expressed through a combination of numbers of order one. At the same time, if the supersymmetry breaking is moderate, as required to solve the quadratic-divergency problem, it may be explained dynamically and related to nonperturbative effects which become important at a characteristic scale of

$$\Lambda \sim \exp\left(-O\left(1/g^2\right)\right) M,$$

where g is some coupling constant. If g is small, then the supersymmetry breaking scale is also small, $\Lambda \ll M$. In a number of realistic models it is possible to get, up to powers of the coupling constants, $v \sim \Lambda$ dynamically (by means of radiative corrections) and to explain therefore the origin of the gauge hierarchy. However, in the MSSM frameworks, there is no place for nonperturbative effects of the required scale: these effects are relevant only for QCD and with $\Lambda \sim \Lambda_{\rm QCD} \ll v$. The dynamical supersymmetry breaking should take place in a new sector, introduced expressly for this



Figure 15: Constraints on the MSSM parameters [134] in one popular scenario (see text). The allowed region is the narrow strip which can be seen in the zoomed panel.

purpose and containing a new strongly coupled gauge theory with its own set of matter fields. No sign of this sector is seen in experiments and one consequently supposes that the interaction between the SM (or MSSM) fields and this sector is rather weak and becomes significant only at high energies, unreachable in the present experiments. This interaction is responsible for the soft terms, that is for mediation of the supersymmetry breaking from the invisible sector to the MSSM sector. One distinguishes the gravity mediation (at Planck energies) and the gauge mediation of supersymmetry breaking. Gravity-mediated and gauge-mediated models have quite different phenomenology.

We see that MSSM, with addition of a sector which breaks supersymmetry dynamically and of a certain interaction between this hidden sector and the observable fields, may explain the origin and stability of the gauge hierarchy, if the masses of superpartners are not very high ($\lesssim$ TeV). Note that the searches for supersymmetry in accelerator experiments put serious constraints on the low-energy supersymmetry. Already the fact that superpartners have not been seen at LEP implied that a significant part of the theoretically allowed MSSM parameter space was excluded experimentally. Subsequent results of Tevatron and especially the first LHC data squeeze the allowed region of parameters significantly, so that for "canonical" supersymmetry, only a very narrow and not fully natural region of possible superpartner mass remains allowed. In Fig. 15, theoretical and experimental (as of summer 2011) constraints on the MSSM parameters are plotted for one rather natural and popular scenario of gravity-mediated supersymmetry breaking. The masses of all scalar superpartners at the $M_{\rm GUT}$ energy scale are equal to m_0 in this scenario, while masses of all fermionic superpartners are $M_{1/2}$. Their ratios to the supersymmetric mixing matrix of the Higgs scalars, μ, are given in the plot. In a scenario which explains the gauge hierarchy, the MSSM parameters and the Z-boson mass should be of the same order; for instance, in the model which corresponds to the illustration, the following relation holds,

$$M_Z^2 \simeq 0.2 m_0^2 + 1.8 M_{1/2}^2 - 2\mu^2.$$

The LHC bound, $M_{1/2} \gtrsim 420$ GeV, results in the requirement of not fully natural

cancellations since $1.8M_{1/2}^2 \gtrsim 40M_Z^2$. Together with the absence of a light Higgs boson discussed in Sec. 4.1, this "little hierarchy" problem makes the approach based on supersymmetry less motivated than it looked some time ago, though there exist variations of supersymmetric models where this dificulty is overcome.

3. The Higgs field may be a pseudo-Goldstone boson. The Goldstone theorem guarantees a massless (even with the account of radiative corrections!) scalar particle for each generator of a broken global symmetry. A weak explicit violation of this symmetry allows to give a small mass to this scalar to get the so-called pseudo-Goldstone boson. The same mechanism results in a low but nonzero mass of some composite particles in a strongly-interacting theory (for instance, of the π meson). A direct application of this approach to the Higgs boson is not possible because the interaction of a pseudo-Goldstone particle with other fields contains derivatives and is very different from the SM interactions. Realistic models of this kind with large coupling constants and with interactions without derivatives, at the same time free from quadratic divergencies, are called the "Little Higgs models" (see e.g. [135] and references therein). Diagram by diagram, the absence of quadratic divergencies occurs due to complicated cancellations of contributions of a number of particles with masses of order TeV, in particular of additional massive scalars. Note that to reconcile a large number of new particles with experimental constraints, in particular with those from the precision electroweak measurements, the model requires significant complications.

4. Composite models: besides the Little Higgs models, a composite Higgs scalar is considered in a number of other constructions, see e.g. [136]. In some rather popular models with composite quarks and leptons, the SM matter fields, together with the Higgs boson (or even without it) represent low-energy degrees of freedom of a strongly coupled theory, like hadrons may be considered as low-energy degrees of freedom of QCD. The mass scales of the theory, v in particular, are determined by the scale Λ at which the running coupling constant of the strongly-coupled theory becomes large, analogously to $\Lambda_{\rm QCD}$. The hierarchy $\Lambda \ll M_{\rm Pl}$ is now determined by the evolution of couplings in the fundamental theory. These models generalize, to some extent, the technicolour models, having more freedom in its construction at the price of even more complications in the quantitative analysis. Note that (at least) in some supersymmetric gauge theories, low-energy degrees of freedom may include also gauge fields, so in principle, one may consider models in which all SM particles are composite (see e.g. [116, 136]). On the other hand, the correspondence between strongly coupled four-dimensional models and weakly-coupled five-dimensional theories (see Sec. 5.3) may open prospects for a quantitative study of composite models. It might even happen that the approaches to the gauge-hierarchy problem, based on assumptions of the extra space dimensions, are equivalent to the approaches which invoke strongly coupled composite models. As in other approaches, to explain the hierarchy, the scale Λ should not exceed significantly the electroweak scale v, so that the LHC constraints on compositeness of quark and leptons (roughly $\Lambda \gtrsim (4 \ldots 5)$ TeV) may again be problematic.

Conclusion. All known scenarios which explain the origin and stability of the gauge hierarchy without extreme fine tuning, predict new particles and/or interactions at the energy scale not far above the electroweak scale. Absence of experimental signs of these particles, especially with the account of the first LHC data, questions the ability of these scenarios to solve the hierarchy problem. If the LHC finds the Higgs scalar but will not confirm predictions of any of the models discussed above, nor will find signs of some other, yet not invented, mechanism, then one would have to reconcider the question of the naturalness of the fine tuning. A principally different position, based on the anthropic principle, is seriously discussed but lays beyond the scope of



Figure 16: Masses of the charged SM fermions. The area of each circle is proportional to the mass of the corrsponding particle.

our consideration.

4.3 The fermion mass hierarchy.

As it has already been pointed out, the SM fermionic fields, quarks and leptons, comprise three generations, that is three sets of particles with identical interactions but with very different masses (see Fig. 16 for a pictorial illustration). The hierarchy of these masses is one of the biggest puzzles of particle physics. Indeed, for instance, the electron ($m_e = 0.511$ MeV), the muon ($m_\mu = 105.7$ MeV) and the tau lepton ($m_\tau = 1777$ MeV) carry identical gauge quantum numbers. For quarks, it is convenient to determine the mass matrix whose diagonal elements determine the masses of the quarks of three generations with identical interactions while combinations of non-diagonal elements provide for the possibility of mixing between generations. The hierarchical structure appears both in the diagonal elements (which differ by orders of magnitude) and in the off-diagonal ones (the mixing is suppressed). In the SM frameworks, neutrino are strictly massless and the mixing of charged leptons is absent, but the same hierarchical structure is seen in the set of masses of charged leptons.

As we have discussed in Sec. 2, the experiments of the past decade not only established confidently the fact of the neutrino oscillations (pointing therefore to nonzero neutrino masses and giving the first laboratory indication to the incompleteness of SM) but also opened the possibility of a quantitative study of neutrino masses and of the mixing in the leptonic sector. It is interesting that the neutrino masses and the leptonic mixings also have the hierarchical structure, but it is very different from the corresponding hierarchy in the quark sector: contrary to the suppressed quark mixings, the leptonic mixing is maximal; the hierarchy of neutrino masses is at the same time moderate. A modern theory which succesfully explains the fermion masses should



Figure 17: A model with extra space dimensions which explains the mass hierarchy.

motivate both hierarchical structures and explain why they are different.

Meanwhile, even without the neutrino sector, the intergeneration mass hierarchy is very difficult to explain. A natural idea is to suppose that there is an extra global symmetry which relates the fermionic generations to each other and which is spontaneously broken; however, this approach is not succesful because it implies the existence of a massless Goldstone boson, the so-called familon, whose parameters are strictly constrained by experiments [6].

A model of fermion masses should explain only the origin of the hierarchy: its stability is provided automatically by the fact that all radiative corrections to the fermion-Higgs Yukawa constants, to which the fermion masses are proportional, depend on the energy logarithmically, that is weakly; this does not, however, make the issue significantly less complicated.

An explanation of the hierarchy may be obtained in a model with extra space dimensions (Fig. 17), in which a single generation of particles in six-dimensional spacetime effectively describes three generations in four dimensions [137, 138]. Each multidimensional fermionic field has three linearly independent solutions which are localized on the four-dimensional hypersurface and have different behaviour close to the brane. Denoting as r, θ the polar coordinates in two extra dimensions and considering the brane at $r = 0$, one gets for the three solutions at $r \to 0$,

$$u_0 \sim \text{const} = r^0 \mathrm{e}^{i0\theta}, \quad u_1 \sim r^1 \mathrm{e}^{i1\theta}, \quad u_2 \sim r^2 \mathrm{e}^{i2\theta}.$$

The Higgs scalar has a vacuum expectation value $v(r)$ which depends on r and is nonzero only in the immediate vicinity of the brane. The effective observable fermion

masses are proportional to the overlap integrals

$$m_i \propto \int dr\, d\theta\, v(r)|u_i|^2(r,\theta)$$

of the coordinate-dependent vacuum expectation value v and extra-dimensional parts of the fermionic wave functions which correspond to the three localized solutions ($i = 0, 1, 2$ enumerates three generation of fermions). One can see from Fig. 17 that the resulting m_i are hierarchically different. Therefore, in this model the mass hierarchy follows from the linear independence of eigenfunctions of the Dirac operator in a particular external field. The same model automatically describes the required structure of neutrino masses and mixings [139]. Presently, this model is the only one known in which the hierarchy of families of both charged fermions and neutrinos are obtained on the common grounds. Note that, contrary to other multidimensional models (e.g. [140]), in this model the number of free parameters is smaller than the number of parameters it describes.

Compared to the hierarchy of masses of particles with identical interactions from different generations, the question of the difference of masses of particles within a generation is much easier. For instance, the difference between masses of the τ lepton and the b and t quarks may be explained by different (because of different quantum numbers) renormalization-group evolution of the Yukawa couplings, so that at the Grand-unification scale these constants are equal while at low energies they are different.

5 Theoretical challenges in the description of hadrons.

5.1 Problems of the perturbative QCD.

In this section, we discuss the question about the practical applicability of the quantum field theory to the description of interactions with large coupling constants, and in particular to the low-energy limit of QCD. It would not be an exaggeration to say that most of the theoretical achievements in the quantum field theory in the past two decades were related to this problem. Before proceeding to the discussion of these achievements, let us note that despite a significant progress, the problem of description of strong interactions at low energies in terms of QCD is not solved, so the development of the corresponding methods remains one of the basic tasks of the quantum field theory.

Recall that QCD, which describes the strong interaction at high energies, is a gauge theory with the gauge group $SU(3)_C$ and $N_f = 6$ fermions, quarks, which transform under its fundamental representation, and the same number of antiquarks transforming under the conjugated representation. A peculiarity of the model is that the asymptotic states, in terms of which the quantum theory is constructed, do not coincide with the fundamental fields in terms of which the Lagrangian is written, that is with fermions (quarks) and gauge bosons (gluons). Contrary, the observable particles do not carry the $SU(3)_C$ quantum numbers (this phenomenon is called confinement). The observable strongly interacting particles are hadrons, whose classification and interactions allow to interpret them as bound states of quarks. At the same time, the theory which describes interaction of quarks, QCD, is unable to calculate properties of these bound states. Intuitively, it seems possible to relate confinement and formation of hadrons with the energy dependence of the QCD gauge coupling constant which grows up with the decrease in energy (that is with the increase in distance; the so-called asymptotic freedom) and becomes large, $\alpha_s \sim 1$, at the scale $\Lambda_{\rm QCD} \sim 150$ MeV: when the distance



Figure 18: Electromagnetic pion formfactor [142]: experimental data versus theoretical calculations, perturbative (QCD, the dashed line) and nonperturbative (full lines representing working models which are not derived from QCD). Up to the energy scale ~ 2 GeV, there are no signs of approaching the perturbative regime.

between quarks is increased, the force between them increases as well, and maybe this force binds them to hadrons. This picture is however not fully consistent because at $\alpha_s \gtrsim 1$, the perturbative expansion stops to work and the true energy dependence of the coupling constant is unknown. Indeed, there exist examples of theories with asymptotic freedom but without confinement [141].

To understand the nature of confinement and to describe properties of hadrons from the first principles (and, in the end, to answer whether QCD is applicable to the description of hadrons), one require the methods of the field theory which do not make use of the expansion in powers of the coupling constants (non-perturbative methods). It is natural to assume (and it was assumed for a long time) that the perturbative QCD has to describe well the physics of strong interactions at characteristic energies above few hundred MeV, because the coupling constant becomes large at ~ 150 MeV. A number of recent experimental results related to the measurement of the form factors of π mesons question the applicability of perturbative methods at considerably higher momentum transfer (a few GeV). In general, formfactors are the coefficients by which the true amplitude of a process with composite or extended particles involved differs from the same amplitude calculated for point-like particles with the same interaction. These coefficients are determined by the internal structure of particles (for instance, by the distribution of the electric charge); their particular form depends on the process considered and on the value of the square of the momentum transfer, Q^2. A full theory describing the interaction which keeps the particles in the bound state should allow for derivation of form factors from the first principles. The results of the experimental determination of formfactors of π mesons related to various processes are given in Figs. 18, 19. One may see that the perturbative QCD experience some difficulties in explaining the experiment at the momentum transfer $\lesssim 4$ GeV.

Approaches to nonperturbative description of QCD may be divided into two classes: (1) calculations in QCD beyond the perturbation theory (the only available method here is the numerical calculation of the functional integral on the lattice) and (2) construction



Figure 19: The transitional form factor of the π meson which describes the process $\pi^0 \to \gamma\gamma$: experimental data [143] versus calculations of perturbative QCD. The QCD predicts the behaviour $Q^2F(Q^2) \sim$ const (the horizonthal full line); at least up to $\sqrt{Q^2} \sim 4$ GeV, the experiment points to $Q^2F(Q^2) \sim (Q^2)^{0.5}$ (the dotted line).

of an effective theory in terms of degrees of freedom which correspond to obsevrable particles. In the latter case the main unsolved question is, as a rule, to justify the connection of the effective theory to QCD. To some extent, a progress in this direction became possible within the concept of dual theories discussed below.

5.2 The lattice results.

The Feynman functional integral is a formally strict approach to the quantization of fields, equivalent to other approaches in the domain of applicability of the perturbation theory. It is natural to suppose that in the nonperturbative domain, this method also reproduces the results which would be obtained within the standard frameworks if the means to get them existed. Numerical calculation of the functional integral is possible in lattice calculations in which the continuous and infinite spacetime is replaced by a finite discrete lattice (see e.g. [144]). In modern calculations, the lattices $32^3 \times 64$, that is 32 points in each of the space coordinates and 64 points in time, are used. For physics applications, it is very important that the gauge invariance may be defined in the lattice theory in a strict way.

One of the first serious achievements of the lattice field theory was a discovery that the lattice model with symmetries and field content of QCD exhibits confinement [145]. Subsequent works allowed to refine which particular field configurations are responsible for confinement; the work on this question continues.

The lattice approach allows to calculate the values of masses and decay constants of hadrons, and in recent years, a significant progress in this direction has been achieved (see Fig. 20). The most precise for today results [146] are obtained for the so-called "2+1" parametrization in which masses of u and s quarks are free parameters, the d-quark mass is assumed to be equal to that of the u quark and the contributions of heavy c, b and t quarks are neglected. Besides these two parameters ($m_u = m_d$ and m_s), there is one more, the physical length which corresponds to a unit step of the lattice. To determine the masses of hadrons, these three parameters should be specified, so in



Figure 20: Results of the lattice calculations of the hadron masses. Masses of π, K and Ω mesons are taken as input parameters. The calculations have been performed in the three-quark approximation, $m_u = m_d \neq m_s$. The histogram gives the experimentally measured values of masses [6], the points (with the error-bar rectangles) represent the results of calculations [146].

real calculations one assumes that the masses of, say, π, K and Ω mesons are known while all other masses and decay constants are expressed through them. One might try to fix masses of heavier particles and to calculate those of the lightest ones, but for a confident calculation of masses of light hadrons a large lattice is required. Currently, the mass of the π meson may be calculated only up to an order of magnitude in this way.

At high temperature, one expects a transition to the state in which quarks cannot be confined in hadrons, that is a phase transition. In reality, these conditions appear in nuclei collisions at high-energy colliders; probably, they also took place in a very early Universe. By means of the lattice methods, the existence of this phase transition has been demonstrated, its temperature has been defined and the dependence of the order of the phase transition from the quark masses has been studied [147, 148].

It is an open theoretical question to prove that the continuum limit of a lattice field theory exists (that is the physical results do not depend on the way in which the lattice size tends to infinity and the lattice step tends to zero) and coincides with QCD. It may happen that this proof is impossible in principle unless one finds an alternative way to work with QCD at strong coupling. However, there exist a series of arguments suggesting that the lattice theory indeed describes QCD (first of all, it is the fact that the lattice calculations reproduce experimental results). At the same time, theoretically, the difference between the lattice and continuum theories is large; for instance, topologically stable in the continuum theory configurations, instantons, which determine the structure of vacuum in nonabelian gauge theories, are not always stable on the lattice; the lattice description of chiral fermions (automatic in a continuum theory) requires complicated constructions etc.

5.3 Dual theories: supersymmetric duality and holography.

In the past two decades, in attempts to relate low-energy models of strong interactions to QCD, theorists created a number of succesful descriptions of dynamics of theories with large coupling constants in terms of other theories, in which the perturbation

theory works. These theories, called dual to each other, have coupling constants g_1 and g_2, for which $g_1 \sim 1/g_2$; the knowledge of the Green functions of one theory allows to calculate, following some known rules, the Green functions of another. Note that the theory dual to QCD has not been constructed up to now.

The simplest example of duality (see e.g. [149]) is a theory of the electromagnetic field with magnetic charges. The Maxwell equations in vacuum are invariant with respect to the exchange of the electric field $\mathbf{E}$ and the magnetic field $\mathbf{B}$:

$$\mathbf{E} \mapsto \mathbf{B}, \quad \mathbf{B} \mapsto -\mathbf{E}. \tag{6}$$

This duality breaks down in the presence of electic charges and currents. It may be restored, however, if one assumes that sources of the other kind exist in Nature, namely magnetic charges and currents which correspond to their motion. The self-consistency of the theory requires the Dirac quantization condition: the unit electic charge e and the unit magnetic charge $\tilde{e}$ have to satisfy the relation $e\tilde{e} = 2\pi$. The charge e is the coupling constant of the usual electrodynamics while the magnetic charge $\tilde{e}$ is the coupling constant of the theory of interaction of magnetic charges which is obtained from electrodynamics by the duality transformation (6). Therefore, the weak coupling of electric charges, $e \ll 1$, corresponds to the strong coupling of magnetic ones, $\tilde{e} = 2\pi/e \gg 1$.

The electromagnetic duality is based on the geometrical properties of abelian gauge fields which cannot be directly transferred to the nonabelian case, which is the most interesting phenomenologically. In a way similar but much more complicated dualities appear in supersymmetric nonabelian gauge theories. The best known one is the "electromagnetic" duality in $SU(2)$ supersymmetric theory with two supercharges ($N = 2$) which is related to the names of Seiberg and Witten [150]. From the particle-physics point of view, this model is a $SU(2)$ gauge theory with scalar and fermionic fields transforming under the adjoint representation of the gauge group, whose interaction is invariant under special symmetry. For this model, the effective theory has been calculated which describes the interaction of light composite particles at low energies and the correspondence has been given between the effective low-energy and fundamental degrees of freedom. Like QCD, the fundamental theory is asymptotically free and is in the strong-coupling regime at low energies; the effective theory describes weakly interacting composite particles.

The success of the Seiberg-Witten model gave rise to a hope that the low-energy effective theory for a nonsupersymmetric gauge model with strong coupling, for instance for QCD, may be obtained from the problem already solved by means of addition of supersymmetry-breaking terms to the lagrangians of both the fundamental and the dual theories. The first step in this direction was to consider $N = 1$ supersymmetric gauge theories. Earlier, starting from mid-1980s, a number of exact results have been obtained in these theories by making use of (gouverned by supersymmetry) analitical properties of the effective action [151]. In contrast with the case of $N = 2$ supersymmetry, this is insufficient for the reconstuction of the full effective theory, but the models dual to supersymmetric gauge theories with different gauge groups and matter content have been suggested [152]. Contrary to the $N = 2$ case, it is impossible to prove the duality here, but the conjecture withstood all checks carried out. Moreover, it has been shown that the addition of small soft breaking terms in the Lagrangians of $N = 1$ theories corresponds to a controllable soft supersymmetry breaking in dual models [153]. Unfortunately, one may prove that with the increase of the supersymmetry-breaking parameters (for instance, when superpartner masses tend to infinity, so the $N = 1$

theory becomes QCD), a phase transition happens and the dual description stops to work, so the straightforward application of this approach to QCD is not possible [154]. Also, it is worth noting that the approach does not allow for a quantitative description of dynamics at intermediate energies, when the coupling constants of dual theories are both large. Nevertheless, these methods themselves, as well as the physics intuition based on their application, have played an important role in the development of other modern approaches to the study of dynamics of strongly-coupled theories.

One of the theoretically most beautiful and practically most prospective approaches to the analysis of dynamics of strong interactions at low and intermediate energies is the so-called holographic approach. Its idea is that the dual theories may be formulated in spacetime of different dimensions, in such a way that, for instance, the four-dimensional dynamics of a theory with large coupling constant is equivalent to the five-dimensional dynamics of another theory which is weakly coupled (in a way similar to the two-dimensional description of a three-dimensional object with a hologram). The best-known realization of this approach is based on the AdS/CFT correspondence [155, 156], a practical realization of the duality between a strongly coupled gauge theory with a four-dimensional conformal invariance (CFT = conformal field theory) and a multidimensional supergravity with weak coupling constant. The four-dimensional conformal symmetry includes the Poincare invariance supplemented by dilatations and inversions. An example of a nontrivial four-dimensional conformal theory with large coupling constant g is the $N = 4$ supersymmetric Yang-Mills theory with the gauge group $SU(N_c)$ which, in the limit $N_c \to \infty$, $g^2 N_c \gg 1$, appears to be dynamically equivalent to a certain supergravity theory living on the ten-dimensional $\mathrm{AdS}_5 \times S_5$ manifold, where AdS_5 is the (4+1)-dimensional space with the anti-de-Sitter metrics (5) and S_5 is the five-dimensional sphere (the S_5 factor is almost irrelevant in applications, hence the name, AdS/CFT correspondense). In the limit considered, these two models are equivalent. To proceed with phenomenological applications, one has to break the conformal invariance. As a result, the theory has less symmetries, so the results proven by making use (direct or indirect) of these symmetries are downgraded to conjectures. Nevertheless, this not fully strict approach (called sometimes AdS/QCD) brings interesting phenomenological results.

An example is provided by a five-dimensional gauge theory determined at a finite interval in the z coordinate of the AdS_5 space (other geometries of the extra dimensions are also considered). For the $SU(2) \times SU(2)$ gauge group and a special matter set one gets the effective theory with QCD symmetries. The series of the Kalutza-Klein states corresponds to the sequence of mesons whose masses and decay constants may therefore be calculated directly in the five-dimensional theory. This approach was succesful; it allows to calculate various physical observables (in particular, the π-meson formfactor discussed above) which agree reasonably with data. A disadvantage of the method is that the duality between QCD and the five-dimensional effective theory is not proven. As a result, the choice of the latter is somewhat arbitrary. An undisputable advantage of this approach is its phenomenological success achieved without a large number of tuning parameters, as well as the possibility to calculate observables for intermediate energies and not only in the zero-energy limit. One may hope that in the future, a low-energy effective theory for QCD might be *derived* in the frameworks of this approach.

6 Conclusions.

The Standard model of particle physics gives an excellent description of almost all data obtained at accelerators already for several decades. At the same time, results of both a number of non-accelerator experiments (neutrino oscillations) and astrophysical observations cannot be explained in the frameworks of SM and undoubtedly point to its incompleteness. A more complete theory, yet to be constructed, should allow for a derivation of the SM parameters and for explanation of their, theoretically not fully natural, values. The main unsolved problem of SM itself is to describe the dynamics of gauge theories at strong coupling which would allow to apply QCD to the description of hadrons at low and intermediate energies.

One may hope that in the next few years, the particle theory will get additional experimental information both from the Large Hadron Collider, a powerful accelerator which is bound to explore the entire range of energies related to the electroweak symmetry breaking, and from numerous experiments of smaller scales (in particular, those studying neutrino oscillations, rare processes etc.) and astrophysical observations. Possibly, this information will allow to construct a succesful extension of the Standard Model already in the coming decade.

This work was born (and grew up) from a review lecture given by the author at the Physics department of the Moscow State University. I am indebted to V. Belokurov who suggested to convert this lecture into a printed text, read the manuscript carefully and discussed many points. I thank V. Rubakov and V. Troitsky for attentive reading of the manuscript and numerous discussions, to M. Vysotsky and M. Chernodub for useful discussions related to particular topics and to A. Strumia for his kind permission to use Fig. 15. The work was supported in part by the RFBR grants 10-02-01406 and 11-02-01528, by the FASI state contract 02.740.11.0244, by the grant of the President of the Russian Federation NS-5525.2010.2 and by the "Dynasty" foundation.

References

[1] Krasnikov N V, Matveev V A *New physics at the Large Hadron Collider* Moscow, Krasand, 2011 (in Russian); Krasnikov N V, Matveev V A *Phys. Usp.* **47** 643 (2004)

[2] Burgess C P, Moore G D *The standard model: A primer,* Cambridge University Press, 2006

[3] Cheng T P, Li L F *Gauge Theory Of Elementary Particle Physics,* Oxford, Clarendon, 1984

[4] Gorbunov D S, Rubakov V A *Introduction to the theory of the early universe: hot big bang theory,* World Scientific, 2011

[5] Kobayashi M *Phys. Usp.* **52** 12 (2009)

[6] Nakamura K *et al.* [Particle Data Group] *J. Phys. G* **37** 075021 (2010)

[7] Giunti C, Kim C W *Fundamentals of Neutrino Physics and Astrophysics,* Oxford University Press 2007

[8] Bilenky S M *Phys. Usp.* **46** 1137 (2003)

[9] Akhmedov E K *Phys. Usp.* **47** 117 (2004)

[10] Kudenko Yu G *Phys. Usp.* **54** 549 (2011)

[11] Evans J J arXiv:1107.3846 [hep-ex]

[12] Pontecorvo B *Sov. Phys. JETP* **6** 429 (1957)

[13] Pontecorvo B *Sov. Phys. JETP* **7** 152 (1957)

[14] Maki Z, Nakagawa M, Sakata S *Prog. Theor. Phys.* **28** 870 (1962)

[15] Pontecorvo B *Sov. Phys. JETP* **26**, 984 (1968)

[16] Gribov V N, Pontecorvo B *Phys. Lett. B* **28** 493 (1969)

[17] Bilenky S M, Pontecorvo B *Sov. J. Nucl. Phys.* **24** 316 (1976)

[18] Bilenky S M, Pontecorvo B *Lett. Nuovo Cim.* **17** 569 (1976)

[19] Eliezer S, Swift A R *Nucl. Phys. B* **105** 45 (1976)

[20] Fritzsch H, Minkowski P *Phys. Lett. B* **62** 72 (1976)

[21] Mikheev S P, Smirnov A Y *Sov. J. Nucl. Phys.* **42** 913 (1985)

[22] Mikheev S P, Smirnov A Y *Nuovo Cim. C* **9** 17 (1986).

[23] Wolfenstein L *Phys. Rev. D* **17** 2369 (1978)

[24] Davis R J, Harmer D S, Hoffman K C *Phys. Rev. Lett.* **20** 1205 (1968)

[25] Hirata K S *et al.* [KAMIOKANDE-II Collaboration] *Phys. Rev. Lett.* **63** 16 (1989)

[26] Abazov A I *et al.* [SAGE Collaboration] *Phys. Rev. Lett.* **67** 3332 (1991)

[27] Hampel W *et al.* [GALLEX Collaboration] *Phys. Lett. B* **447** 127 (1999)

[28] Ashie Y *et al.* [Super-Kamiokande Collaboration] *Phys. Rev. Lett.* **93** 101801 (2004)

[29] Hirata K S *et al.* [KAMIOKANDE-II Collaboration] *Phys. Lett. B* **205** 416 (1988)

[30] Casper D *et al. Phys. Rev. Lett.* **66** 2561 (1991)

[31] Allison W W M *et al. Phys. Lett. B* **391** 491 (1997)

[32] Ambrosio M *et al.* [MACRO Collaboration] *Phys. Lett. B* **434** 451 (1998)

[33] Fukuda Y *et al.* [Super-Kamiokande Collaboration] *Phys. Rev. Lett.* **82** 2644 (1999)

[34] Ahmad Q R *et al.* [SNO Collaboration] *Phys. Rev. Lett.* **89** 011301 (2002)

[35] Abe S *et al.* [KamLAND Collaboration] *Phys. Rev. Lett.* **100** 221803 (2008)

[36] Aharmim B *et al.* [SNO Collaboration] arXiv:1109.0763

[37] Bellini G *et al.* [Borexino Collaboration] arXiv:1104.1816 [hep-ex].

[38] Ashie Y *et al.* [Super-Kamiokande Collaboration] *Phys. Rev. D* **71** 112005 (2005)

[39] Takeuchi Y *et al.* [Super-Kamiokande Collaboration] talk at Neutrino-2010, Athens, 14-19 June 2010.

[40] Ahn M H *et al.* [K2K Collaboration] *Phys. Rev. D* **74** 072003 (2006)

[41] Michael D G *et al.* [MINOS Collaboration] *Phys. Rev. Lett.* **97** 191801 (2006)

[42] Adamson P *et al.* [The MINOS Collaboration] *Phys. Rev. Lett.* **106** 181801 (2011)

[43] Agafonova N *et al.* [OPERA Collaboration] *Phys. Lett. B* **691** 138 (2010)

[44] Fogli G L *et al., Phys. Rev. Lett.* **101** 141801 (2008)

[45] Abe K *et al.* [T2K Collaboration] *Phys. Rev. Lett.* **107** 041801 (2011)

[46] Adamson P *et al.* [MINOS Collaboration] arXiv:1108.0015 [hep-ex]

[47] Fogli G L *et al.,* arXiv:1106.6028 [hep-ph]

[48] Aguilar A *et al.* [LSND Collaboration] *Phys. Rev. D* **64** 112007 (2001)

[49] Church E D *et al., Phys. Rev. D* **66** 013001 (2002)

[50] Aguilar-Arevalo A A *et al.* [The MiniBooNE Collaboration] *Phys. Rev. Lett.* **105** 181801 (2010)

[51] Djurcic Z, talk at *13th International Workshop on Neutrino Factories, Super Beams and Beta Beams*, Geneva, 1–6 August 2011

[52] Mention G *et al. Phys. Rev. D* **83** 073006 (2011)

[53] Aguilar-Arevalo A A *et al.* [MiniBooNE Collaboration] *Phys. Rev. Lett.* **103** 111801 (2009)

[54] Thomas J, talk at *Lepton-Photon 2011*, Mumbai, 22–27 August 2011

[55] Abe K *et al.* [Kamiokande Collaboration] arXiv:1109.1621

[56] Anselmann P *et al.* [GALLEX Collaboration.] *Phys. Lett. B* **342** 440 (1995); Kaether F *et al. Phys. Lett. B* **685** 47 (2010)

[57] Abdurashitov D N *et al.* [SAGE Collaboration] *Phys. Rev. Lett.* **77** 4708 (1996); Abdurashitov D N *et al.* [SAGE Collaboration] *Phys. Rev. C* **73** 045805 (2006)

[58] Giunti C, Laveder M *Phys. Rev. C* **83** 065504 (2011)

[59] Giunti C, Laveder M *Phys. Rev. D* **82** 113009 (2010)

[60] Aguilar-Arevalo A A *et al.* [MiniBooNE Collaboration] *Phys. Rev. Lett.* **102** 101802 (2009)

[61] Lobashev V M *et al. Phys. Lett.* B **460** 227 (1999)

[62] Aguilar-Arevalo A A *et al.* [MiniBooNE Collaboration] [arXiv:1109.3480 [hep-ex]]

[63] Adam T *et al.* [OPERA Collaboration] arXiv:1109.4897 [hep-ex]

[64] Strumia A *Phys. Lett. B* **539** 91 (2002)

[65] Maltoni M *et al. Nucl. Phys. B* **643** 321 (2002)

[66] Akhmedov E, Schwetz T *JHEP* **1010** 115 (2010)

[67] Murayama H, Yanagida T *Phys. Lett. B* **520** 263 (2001)

[68] Bogolyubov N N, Shirkov D V *Introduction to the theory of quantized fields,* Intersci. Monogr. Phys. Astron. **3** 1 (1959)

[69] Tsukerman I S *Phys. Usp.* **48** 825 (2005); Tsukerman I S arXiv:1006.4989 [hep-ph]

[70] Diaz J S, Kostelecky A arXiv:1108.1799 [hep-ph]

[71] Engelhardt N, Nelson A E, Walsh J R *Phys. Rev. D* **81** 113001 (2010)

[72] Kopp J, Machado P A N, Parke S J *Phys. Rev. D* **82** 113002 (2010)

[73] Schwetz T arXiv:0805.2234 [hep-ph]

[74] Yasuda O arXiv:1012.3478 [hep-ph]

[75] Aseev V N *et al.*, arXiv:1108.5034 [hep-ex]

[76] Kraus C *et al. Eur. Phys. J. C* **40** 447 (2005)

[77] Hannestad S *et al. JCAP* **1008** 001 (2010)

[78] Gorbunov D S, Rubakov V A *Introduction to the theory of the early universe, Cosmological perturbations and inflationalry theory*, World Scientific, 2011

[79] Rubakov V A *Phys. Usp.* **42** 1193 (1999)

[80] Rubakov V A *Phys. Usp.* **54** 633 (2011)

[81] Sajharov A D *JETP Lett.*, **5** 24 (1967)

[82] Rubakov V A, Shaposhnikov M E *Phys. Usp.* **39** 461 (1996)

[83] Rubin V C, Thonnard N, Ford W K *Astrophys. J.* **238** 471 (1980)

[84] Begeman K G *Astron. Astrophys.* **223** 47 (1989)

[85] The digitized sky survey (DSS), in [88].

[86] Zwicky F *Astrophys. J.* **86** 217 (1937)

[87] Limousin M *et al. Astrophys. J.* **668** 643 (2007); http://www.dark-cosmology.dk

[88] The Multimission Archive at the Space Telescope Science Institute (MAST), http://archive.stsci.edu/ . STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555.

[89] The Chandra Data Archive (CDA), http://asc.harvard.edu/cda/ .

[90] Clowe D *et al.* *Astrophys. J.* **648** L109 (2006)

[91] Bradac M *et al.* *Astrophys. J.* **652** 937 (2006)

[92] Alcock C *et al.* [MACHO Collaboration] *Astrophys. J.* **542** 281 (2000); Tisserand P *et al.* [EROS-2 Collaboration] *Astron. Astrophys.* **469** 387 (2007)

[93] Riess A G *et al.* [Supernova Search Team Collaboration] *Astron. J.* **116** (1998) 1009

[94] Perlmutter S *et al.* [Supernova Cosmology Project Collaboration] *Astrophys. J.* **517** 565 (1999)

[95] Perlmutter S *Physics Today* **56** (4) 53 (2003)

[96] Riess A G, Press W H, Kirshner R P *Astrophys. J.* **473** 88 (1996)

[97] Perlmutter S *et al.* [Supernova Cosmology Project Collaboration] Astrophys. J. **483** (1997) 565

[98] Amanullah R *et al. Astrophys. J.* **716** 712 (2010); data for Fig. 11 are taken from http://supernova.lbl.gov/Union/

[99] Jullo E *et al. Science* **329** 924 (2010)

[100] Komatsu E *et al.* [WMAP Collaboration] *Astrophys. J. Suppl.* **192** 18 (2011)

[101] Marinoni C, Buzzi A *Nature* **468** 539 (2010)

[102] Khoury J, Weltman A *Phys. Rev. D* **69** 044026 (2004)

[103] Linde A D *Particle physics and inflationary cosmology,* Harwood Academic Publishers, 1990

[104] Barate R *et al.* [LEP Working Group for Higgs boson searches, ALEPH, DELPHI, L3 and OPAL Collaborations] *Phys. Lett. B* **565** 61 (2003)

[105] Sharma V *et al.* [CMS Collaboration], talk at *Lepton-Photon 2011*, Mumbai, 22–27 August 2011

[106] Nisati A *et al.* [ATLAS Collaboration], talk at *Lepton-Photon 2011*, Mumbai, 22–27 August 2011

[107] Verzocchi M *et al.* [CDF and D0 Collaborations], talk at *Lepton-Photon 2011*, Mumbai, 22–27 August 2011

[108] Baak M *et al.* arXiv:1107.0975

[109] Grojean C *Phys. Usp.* **50** 1 (2007)

[110] Lane K arXiv:hep-ph/0202255

[111] Manton N S *Nucl. Phys. B* **158** 141 (1979)

[112] Csaki *et al. Phys. Rev. D* **69** 055006 (2004)

[113] Rubakov V A *Phys. Usp.* **50** 390 (2007)

[114] Atwood D, Gupta S K, Soni A arXiv:1104.3874 [hep-ph]

[115] Ghilencea D, Lanzagorta M, Ross G G *Phys. Lett. B* **415** 253 (1997)

[116] Rubakov V A, Troitsky S V arXiv:hep-ph/0001213

[117] Rubakov V A *Phys. Usp.* **46** 211 (2003)

[118] Kaluza T *Sitzungsber. Preuss. Akad. Wiss. Berlin*, Math.-Phys. Kl. (1) 966 (1921)

[119] Klein O *Z. Phys.* **37** 895 (1926)

[120] Akama K *Lecture Notes Phys.* **176** 267 (1983)

[121] Rubakov V A, Shaposhnikov M E *Phys. Lett. B* **125** 136 (1983)

[122] Visser M *Phys. Lett. B* **159** 22 (1985)

[123] von Klitzing K, Nobel lecture (1985); Fu L, Kane C L *Phys. Rev. Lett.* **100** 096407 (2008); Ghaemi P, Wilczek F arXiv:0709.2626; Bergman D L, Le Hur K *Phys. Rev. B* **79** 184520 (2009); Volovik G E *The Universe in a helium droplet*, *Int. Ser. Monogr. Phys.* **117** (2006)

[124] Arkani-Hamed N, Dimopoulos S, Dvali G *Phys. Lett. B* **429** 263 (1998)

[125] Kapner D J *et al. Phys. Rev. Lett.* **98** 021101 (2007)

[126] Dvali G, Shifman M *Phys. Lett. B* **396** 64 (1997)

[127] Rubakov V A, Shaposhnikov M E *Phys. Lett. B* **125** 139 (1983)

[128] Gogberashvili M *Mod. Phys. Lett. A* **14** 2025 (1999)

[129] Randall L, Sundrum R *Phys. Rev. Lett.* **83** 3370 (1999)

[130] Oda I *Phys. Lett. B* **496** 113 (2000)

[131] Vysotsky M I, Nevzorov R B *Phys. Usp.* **44** 919 (2001)

[132] Gorbunov D S, Dubovsky S L, Troitsky S V *Phys. Usp.* **42** 623 (1999)

[133] Kazakov D I arXiv:hep-ph/0012288

[134] Strumia A *JHEP* **1104** 073 (2011)

[135] Schmaltz M, Tucker-Smith D *Ann. Rev. Nucl. Part. Sci.* **55** 229 (2005)

[136] Gherghetta T arXiv:1008.2570 [hep-ph].

[137] Libanov M, Troitsky S *Nucl. Phys. B* **599** 319 (2001)

[138] Frere J-M, Libanov M, Troitsky S *Phys. Lett. B* **512** 169 (2001)

[139] Frere J-M, Libanov M, Ling F S *JHEP* **1009** 081 (2010)

[140] Dvali G R, Shifman M A, *Phys. Lett.* **B475** 295 (2000)

[141] Iwasaki Y *et al. Phys. Rev. Lett.* **69** 21 (1992)

[142] Krutov A F, Troitsky V E, Tsirova N A *Phys. Rev. C* **80** 055210 (2009)

[143] Aubert B *et al.* [The BABAR Collaboration] *Phys. Rev. D* **80** 052002 (2009)

[144] Di Giacomo A *Lattice gauge theory*, in: *Encyclopedia of Mathematical Physics*, Academic Press, Oxford (2006)

[145] Creutz M *Phys. Rev. D* **21** 2308 (1980)

[146] Aoki S *et al.* [PACS-CS Collaboration] *Phys. Rev. D* **81** 074503 (2010)

[147] Kajantie K, Montonen C, Pietarinen E *Z. Phys. C* **9** 253 (1981)

[148] Aoki Y *et al.* Nature **443** 675 (2006)

[149] Tsun T S *Electric–magnetic duality*, in: *Encyclopedia of Mathematical Physics*, Academic Press, Oxford (2006)

[150] Seiberg N, Witten E *Nucl. Phys. B* **426** 19 (1994) [Erratum-ibid. **430** 486 (1994)]

[151] Affleck I, Dine M, Seiberg N *Nucl. Phys. B* **241** 493 (1984)

[152] Seiberg N *Nucl. Phys. B* **435** 129 (1995)

[153] Evans N J, Hsu S D H, Schwetz M *Phys. Lett. B* **355** 475 (1995)

[154] Aharony O *et al. Phys. Rev. D* **52** 6157 (1995)

[155] Maldacena J M *Adv. Theor. Math. Phys.* **2** 231 (1998) [*Int. J. Theor. Phys.* **38** 1113 (1999)]

[156] Witten E *Adv. Theor. Math. Phys.* **2** 253 (1998)